# QUADRATIC SEGRE INDICES

#### FELIPE ESPREAFICO, STEPHEN MCKEAN, AND SABRINA PAULI

ABSTRACT. We prove that the local Euler class of a line on a degree 2n-1 hypersurface in projective n + 1 space is given by the product of indices of Segre involutions. Segre involutions and their associated indices were first defined by Finashin and Kharlamov over the reals. Our result is valid over any field of characteristic not 2 and gives an infinite family of problems in enriched enumerative geometry with a shared geometric interpretation for the local type.

#### 1. INTRODUCTION

We begin with a brief statement of our main result. We then give the motivation for our result, as well as the relevant terminology, in Section 1.1. Sufficiently motivated readers who are tired of history may skip to Section 1.2, where we summarize the ideas behind our proofs.

In short, we give a geometric interpretation for the enumerative weight of a line on a general degree 2n - 1 hypersurface in  $\mathbb{P}^{n+1}$  over an arbitrary field of characteristic not 2, generalizing work of Kass–Wickelgren in the n = 2 case [KW21] and Pauli in the n = 3 case [Pau22]. Our geometric interpretation is directly inspired by a construction of Finashin and Kharlamov [FK21].

Setup 1.1. Let  $X \subset \mathbb{P}_k^{n+1}$  be a general hypersurface of degree 2n-1 over a field k. Given a line  $\ell \subset X$  with field of definition  $k(\ell)$ , there is an associated rational curve  $\mathcal{G}(\ell) \subset \mathbb{P}_k^{n-1}$  of degree 2n-2. Over  $\overline{k}$  there are exactly  $\binom{n}{2}$  planes of dimension n-3 meeting  $\mathcal{G}(\ell)$  in 2n-4 points. Each such (2n-4)-secant plane S, whose field of definition we denote k(S), determines an involution  $i_S : \ell_{k(S)} \to \ell_{k(S)}$ . The fixed points of  $i_S$  are defined over  $k(S)(\sqrt{\alpha_S})$  for some  $\alpha_S \in k(S)^{\times}/(k(S)^{\times})^2$ .

Let  $\mathrm{GW}(k)$  denote the Grothendieck–Witt ring of virtual quadratic forms over k. Given  $a \in k^{\times}$ , let  $\langle a \rangle \in \mathrm{GW}(k)$  represent the isomorphism class of the bilinear form  $[(x, y) \mapsto axy]$ . Given a field extension L/k, let  $\mathrm{N}_{L/k} : L \to k$  and  $\mathrm{Tr}_{L/k} : L \to k$  denote the field norm and field trace, respectively.

We define the Segre index of  $\ell \subset X$  to be

$$\operatorname{seg}(X, \ell) := \operatorname{Tr}_{k(\ell)/k} \left\langle \prod_{S} \operatorname{N}_{k(S)/k(\ell)}(\alpha_{S}) \right\rangle \in \operatorname{GW}(k),$$

<sup>2020</sup> Mathematics Subject Classification. Primary: 14N15. Secondary: 14G27.

where this product ranges over the (n-3)-planes that are (2n-4)-secant to  $\mathcal{G}(\ell)$ . Note that  $k(\ell)/k$  is separable for general X by e.g. [EH16, Theorem 6.34], so the field trace here is non-degenerate.

**Theorem 1.2** (Main Theorem). Let  $X = \mathbb{V}(F) \subset \mathbb{P}_k^{n+1}$  be a general hypersurface of degree 2n-1 over a field k. Let

$$\sigma_F : \mathbb{G}(1, n+1) \to \operatorname{Sym}^{2n-1}(\mathcal{S}^{\vee})$$

denote the section determined by F, where S is the tautological bundle on the Grassmannian of lines in  $\mathbb{P}^{n+1}$ . Let  $\operatorname{ind}_{\ell} \sigma_F \in \operatorname{GW}(k)$  denote the local index of  $\sigma_F$  at a line  $\ell \subset X$ . Then

$$\operatorname{ind}_{\ell} \sigma_F = \operatorname{seg}(X, \ell).$$

Theorem 1.2 allows us to immediately deduce a quadratically enriched count of lines on a degree 2n - 1 hypersurface in  $\mathbb{P}^{n+1}$ . This is because the Euler number underlying the total count is completely determined by the (signed) real and complex counts [Lev19, Example 8.2], [BW23, Corollary 6.9], the real count is (2n - 1)!! [FK12, OT14], and the complex count is computable by Schubert calculus.

**Theorem 1.3.** Let  $X \subset \mathbb{P}_k^{n+1}$  be a general hypersurface of degree 2n-1 over a field k. Let c(n) denote the top Chern number of  $\operatorname{Sym}^{2n-1}(\mathcal{S}^{\vee}) \to \mathbb{G}(1, n+1)$ . Then we have an equality

$$\sum_{\ell \subset X} \operatorname{seg}(X, \ell) = (2n-1)!! \langle 1 \rangle + \frac{c(n) - (2n-1)!!}{2} \mathbb{H}$$

in GW(k).

These theorems provide an infinite family of enriched enumerative problems with a shared geometric interpretation for their local indices. In the language of the *geometricity question* [McK22, Appendix C], these "lines on hypersurfaces" problems belong to the same phylum of enumerative problems, as one would expect.

1.1. **Motivation.** Famously, there are 27 lines on every smooth complex cubic surface [Cay49]. A modern method for calculating the number 27 is as the top Chern number of the bundle

$$\operatorname{Sym}^{3}(\mathcal{S}^{\vee}) \to \mathbb{G}(1,3),$$

where  $S \to \mathbb{G}(1,3)$  is the tautological bundle on the Grassmannian of projective lines in  $\mathbb{P}^3$ . More generally, one can prove that there are finitely many lines on a generic degree 2n-1 hypersurface in  $\mathbb{P}^{n+1}$ , all of which are reduced. The top Chern number of

$$\operatorname{Sym}^{2n-1}(\mathcal{S}^{\vee}) \to \mathbb{G}(1, n+1)$$

can then be computed via Schubert calculus, giving the 2875 lines on a quintic threefold or, for those impressed by large numbers, the 1,192,221,463,356,102,320,754,899 lines on a novemdecic tenfold (see e.g. [EH16, §6.5]).

Lines on hypersurfaces generally make for good conversation with non-mathematicians, such as relatives or university administrators — most people can appreciate the visual



FIGURE 1. Segre involution on a cubic surface

beauty of a cubic surface with its configuration of lines, and administrators tend to be impressed by large numbers. However, as with all online image searches, one should be careful when pulling up a picture of a cubic surface and its lines at a family party. This is because any such picture will actually depict a *real* cubic surface, which may have fewer than 27 *real* lines.

1.1.1. The real story. Schläffi proved that every smooth real cubic surface has 3, 7, 15, or 27 real lines [Sch58]. Nearly a century later, Segre constructed an involution on each real line on a cubic surface [Seg42], which we now describe. Let  $\ell$  be a real line on a real cubic surface X. For each point  $p \in \ell$ , the intersection  $T_pX \cap X$  is the union of  $\ell$  and a residual conic  $Q_p \subset T_pX$ . The intersection  $Q_p \cap \ell$  consists of p and another point, say q. The Segre involution swaps p and q (see Figure 1).

Define hyperbolic lines as those whose Segre involution has real fixed points and elliptic lines as those whose Segre involution has complex fixed points. For each topological type of real cubic surface, Segre computed the number of hyperbolic and elliptic lines contained therein. These computations imply the striking formula

(1.1) 
$$\#\{\text{hyperbolic lines}\} - \#\{\text{elliptic lines}\} = 3,$$

although this seems to have gone unnoticed until it was observed by Finashin–Kharlamov [FK12] and Okonek–Teleman [OT14] about 70 years later. Equation 1.1 should be viewed as the correct analog of the 27 lines on a cubic surface over  $\mathbb{C}$ . While the total number of real lines on a cubic surface over  $\mathbb{R}$  depends on the choice of cubic surface, the *signed* count of lines given in Equation 1.1 is independent of the choice of cubic surface.

Inspired by Equation 1.1, Finashin and Kharlamov set out to give a signed count of real lines on degree 2n-1 hypersurfaces in  $\mathbb{RP}^{n+1}$ . They proved that the overall signed count is equal to (2n-1)!! [FK12], which arises as the Euler number of the real vector bundle  $\operatorname{Sym}^{2n-1}(\mathcal{S}^{\vee}) \to \mathbb{G}(1, n+1)$ . See also [OT14, Sol06] for related results. To complete the signed count of real lines on hypersurfaces, there also needs to be a geometric determination of the type of a line, analogous to Segre involutions determining whether a line on a cubic surface is hyperbolic or elliptic. Finashin and Kharlamov gave two such geometric interpretations (and proved that they are equivalent) in [FK21]: Welschinger weights and Segre indices. We will only discuss Segre indices, as these are the interpretation relevant for our article. Given a smooth hypersurface  $X \subset \mathbb{RP}^{n+1}$ , there is a *Gauß map* 

$$\mathcal{G}: X \to \mathbb{RP}^{n+1}$$
$$p \mapsto T_p X,$$

where the target  $\mathbb{RP}^{n+1}$  is the dual projective space parameterizing codimension 1 hyperplanes in  $\mathbb{RP}^{n+1}$ . If we restrict  $\mathcal{G}$  to a line  $\ell \subset X$ , then the tangent plane  $\mathcal{G}(p) := T_p X$  contains  $\ell$  for each  $p \in \ell$ . The flag variety of codimension 1 hyperplanes in  $\mathbb{RP}^{n+1}$  containing a given line (in this case,  $\ell$ ) is isomorphic to  $\mathbb{RP}^{n-1}$ . We thus obtain a restricted Gauß map

$$\mathcal{G}|_{\ell}: \ell \to \mathbb{RP}^{n-1}.$$

The image  $\mathcal{G}(\ell) \subset \mathbb{RP}^{n-1}$  is a rational curve of degree 2n-2, which we refer to as the *Gauß curve* of  $\ell$ . For  $n \geq 3$ , a generic degree 2n-2 rational curve in  $\mathbb{RP}^{n-1}$  has a finite number of (n-3)-dimensional (2n-4)-secants; in fact, there are  $\binom{n}{2}$  such secants defined over  $\mathbb{C}$ .

Each secant defined over  $\mathbb{R}$  determines a Segre involution as follows. Let S be an (n-3)-plane meeting  $\mathcal{G}(\ell)$  in 2n-4 points defined over  $\mathbb{R}$ . There is a pencil of (n-2)-dimensional (i.e. codimension 1 in  $\mathbb{P}^{n-1}$ ) hyperplanes containing S. Each (n-2)-plane in this pencil meets the Gauß curve  $\mathcal{G}(\ell)$  in 2n-2 points, by Bézout's theorem: the 2n-4 secant points given by  $S \cap \mathcal{G}(\ell)$ , and another pair of points. The Segre involution associated to S swaps this residual pair of points, as depicted in Figure 2.<sup>1</sup>

The Segre index of  $\ell$ , which we denote  $\operatorname{seg}(X, \ell)$ , is a product of indices associated to the Segre involutions on  $\mathcal{G}(\ell)$  corresponding to the secant defined over  $\mathbb{R}$ . This Gauß curve is the embedding of a conic  $Q_{\ell} \subset \mathbb{RP}^2$ , and the Segre involution associated to a secant S pulls back to an involution on  $Q_{\ell}$ . Define the index of this involution, denoted  $\operatorname{seg}_S(X, \ell)$ , to be +1 if its fixed points are real and -1 if its fixed points are complex. Then

$$\operatorname{seg}(X,\ell) := \prod_{\operatorname{real secants } S} \operatorname{seg}_S(X,\ell),$$

and Finashin and Kharlamov proved that

(1.2) 
$$\sum_{\ell \subset X} \sec(X, \ell) = (2n - 1)!!$$

for a generic degree 2n - 1 hypersurface in  $\mathbb{RP}^{n+1}$ . Note that Equation 1.1 is the n = 2 case of Equation 1.2. For n = 2, the Segre index agrees with the convention that assigns +1 to hyperbolic lines and -1 to elliptic lines on a cubic surface.

<sup>&</sup>lt;sup>1</sup>The plane quartic in Figure 2A is the well-known ampersand curve. To construct a rational space sextic curve (like the one in Figure 2B) and its six quadrisecants, it helps to use a remarkable theorem of Dye stating that such quadrisecants form half of a double six of lines on a cubic surface containing the sextic [Dye97]. See Remark 3.12 for more details.



FIGURE 2. Segre involutions

1.1.2. The arithmetic story. Our goal (stated previously as Theorem 1.3) is to prove an analog of Equation 1.2 over an arbitrary base field. The framework for such a theorem is the *enriched enumerative geometry* program, in which Kass and Wickelgren's count of lines on cubic surfaces is an early seminal result [KW21].

Let k be a field. We assume that char  $k \neq 2$ , as we want involutions to behave well over k. Let  $\mathrm{GW}(k)$  denote the Grothendieck–Witt ring of isomorphism classes of symmetric nondegenerate bilinear forms. This ring is generated by classes of the form  $\langle a \rangle = [(x, y) \mapsto axy]$  for  $a \in k^{\times}$ . Kass and Wickelgren defined an Euler "number"  $e(\mathrm{Sym}^3(\mathcal{S}^{\vee})) \in \mathrm{GW}(k)$ that satisfies a Poincaré–Hopf theorem: given a smooth cubic surface X over k, the associated section  $\sigma_X : \mathbb{G}(1,3) \to \mathrm{Sym}^3(\mathcal{S}^{\vee})$  has local indices  $\mathrm{ind}_{\ell} \sigma_X \in \mathrm{GW}(k)$  such that

(1.3) 
$$\sum_{\ell \subset X} \operatorname{ind}_{\ell} \sigma_X = e(\operatorname{Sym}^3(\mathcal{S}^{\vee})).$$

In order to turn Equation 1.3 into an enumerative theorem, one needs to compute the Euler number on the right hand side and give a geometric interpretation of the local indices on the left hand side. Kass and Wickelgren computed

$$e(\operatorname{Sym}^{3}(\mathcal{S}^{\vee})) = 15\langle 1 \rangle + 12\langle -1 \rangle$$

and proved that  $\operatorname{ind}_{\ell} \sigma_X$  is determined by the Segre involution associated to  $\ell$ . Just as the Segre involution of a real line has either real or complex fixed points, the fixed points of the Segre involution on a line with field of definition  $k(\ell)$  are defined over  $k(\ell)(\sqrt{\alpha_{\ell}})$  for some  $\alpha_{\ell} \in k(\ell)^{\times}/(k(\ell)^{\times})^2$ . Kass and Wickelgren proved that

$$\operatorname{ind}_{\ell} \sigma_X = \operatorname{Tr}_{k(\ell)/k} \langle \alpha_{\ell} \rangle \in \operatorname{GW}(k),$$

giving the desired geometric description of the relevant local indices. Altogether, we get the count

(1.4) 
$$\sum_{\ell \subset X} \operatorname{Tr}_{k(\ell)/k} \langle \alpha_{\ell} \rangle = 15 \langle 1 \rangle + 12 \langle -1 \rangle.$$

One of the key features of Equation 1.4 is that it generalizes both the complex and real counts of lines on cubic surfaces. The rank of Equation 1.4 states that

$$\sum_{\ell \subset X_{\overline{k}}} 1 = 27$$

counting 27 geometric lines on every smooth cubic surface. When k admits a real embedding, the signature of Equation 1.4 exactly recovers Equation 1.1, as hyperbolic and elliptic lines respectively satisfy  $\alpha_{\ell} = +1$  and  $\alpha_{\ell} = -1$ , while the signature of  $\text{Tr}_{\mathbb{C}/\mathbb{R}} \langle \alpha_{\ell} \rangle$ is 0 for any complex line.

Lines on quintic threefolds over general fields were considered by Levine and Pauli. Levine [Lev19, Example 8.3] computed

$$e(\operatorname{Sym}^5(\mathcal{S}^{\vee})) = 1445\langle 1 \rangle + 1430\langle -1 \rangle$$

while Pauli [Pau22] provided a geometric interpretation of the local indices contributing to this Euler number. The geometric interpretation is a quadratic enrichment of (the n = 3 case of) the Segre index described in Section 1.1.1. In this case, the Gauß curve  $\mathcal{G}(\ell)$  associated to a line  $\ell$  is a plane quartic, and geometrically there are  $\binom{3}{2} = 3$  planes of dimension 3 - 3 = 0 that meet this quartic with order  $2 \cdot 3 - 4 = 2$ . In other words,  $\mathcal{G}(\ell)$  is a plane quartic with 3 nodes. The Segre involution

$$i_{\nu}: \mathcal{G}(\ell) \to \mathcal{G}(\ell)$$

associated to a node  $\nu \in \mathcal{G}(\ell)$  is given by taking a pencil of lines through  $\nu$ , intersecting each line L with  $\mathcal{G}(\ell)$ , and swapping the pair of points in  $\mathcal{G}(\ell) \cap L - \nu$  (see Figure 2A, where  $S = \nu$ ). The fixed points of  $i_{\nu}$  are defined over  $k(\nu)(\sqrt{\alpha_{\nu}})$  for some  $\alpha_{\nu} \in k(\nu)^{\times}/(k(\nu)^{\times})^2$ , and the Segre index of  $\ell$  is defined as

$$\operatorname{seg}(X,\ell) := \operatorname{Tr}_{k(\ell)/k} \left\langle \prod_{\nu} \operatorname{N}_{k(\nu)/k(\ell)}(\alpha_{\nu}) \right\rangle.$$

**Remark 1.4.** While we have described the Segre involution associated to a node  $\nu$  as an involution of the form  $i_{\nu} : \mathcal{G}(\ell) \to \mathcal{G}(\ell)$ , one can equivalently consider the pullback of  $i_{\nu}$  to an involution  $\ell$ . We will prove that the fixed points of the involutions on  $\mathcal{G}(\ell)$  and  $\ell$ have isomorphic residue fields. This will imply that  $\operatorname{seg}(X, \ell)$  is independent of whether we think of the involution occurring on  $\mathcal{G}(\ell)$  or  $\ell$ , so we will generally not distinguish between these two perspectives.

Already at the time of [Pau22], it was expected that a similar description of the local index in terms of Segre involutions should hold in the general case of lines on degree 2n-1 hypersurfaces in  $\mathbb{P}^{n+1}$ . However, the proof in [Pau22] utilizes the fact that for a

general rational plane quartic, its trio of nodes lie in general position (i.e. not on a line). It is not true that the (n-3)-dimensional (2n-4)-secants to a general degree 2n-2 space curve in  $\mathbb{P}^{n-1}$  are in general position when n > 3, since  $\binom{n}{2} > \dim \mathbb{G}(n-3, n-1)+1$  in this range. Furthermore, the proofs in [FK21] rely on wall-crossing arguments, which do not readily generalize to other fields.

As we have claimed in Theorem 1.2, the quadratic Segre index of a line indeed computes the local index contributing to the Euler class  $e(\text{Sym}^{2n-1}(\mathcal{S}^{\vee}))$ . In the next subsection, we will outline the proof ideas that allowed us to sidestep the difficulties that arise when trying to generalize from  $\mathbb{R}$  to k or from n = 3 to n > 3.

1.2. Ideas behind the proof. As previously mentioned, we use a construction of Finashin and Kharlamov [FK21, Section 6] to relate lines on hypersurfaces to involutions arising from the secants of their associated Gauß curves, and equivalently to involutions associated to plane conics avoiding an auxiliary set of points. This construction allows us to generalize beyond the cases of cubic surfaces [KW21] and quintic threefolds [Pau22].

In order to show that the local index of a line is equal to the product of Segre indices associated to the Gauß curve or plane conic, we essentially need to show that two determinants are equal up to squares<sup>2</sup>. Over  $\mathbb{R}$ , this boils down to showing that two real numbers have the same sign; this is where Finashin–Kharlamov utilize a wall-crossing argument. Over arbitrary fields, we instead will show that these two determinants are given by regular maps with the same zero locus. This step more or less falls out of the geometry, which will allow us to express these regular maps in terms of resultants. The primary difficulties arise from the fact that conics over non-closed fields need not have rational points. We overcome this difficulty by modifying Finashin and Kharlamov's construction and working with *parameterized* conics.

The final step is to show that the relevant regular maps agree at a point outside of their zero locus. This is done with an elementary, although somewhat lengthy, argument involving only basic linear algebra.

1.3. Outline. Here is a quick outline of the article.

- In Section 2, we discuss the bundle  $\operatorname{Sym}^{2n-1} \mathcal{S}^{\vee} \to \mathbb{G}(1, n+1)$ , the section induced by a hypersurface, our choice of coordinates on  $\mathbb{G}(1, n+1)$ , and our local trivializations of  $\operatorname{Sym}^{2n-1} \mathcal{S}^{\vee}$ . We then compute the local index and give examples discussing the cases of cubic surfaces and quintic threefolds.
- In Section 3, we discuss Finashin and Kharlamov's conic models for rational space curves. This includes a dictionary translating between Gauß curves with their secants and plane conics with their associated loci of points. We also prove that the index of a Gauß curve can be computed in terms of a conic model.

<sup>&</sup>lt;sup>2</sup>We say that two elements x, y of a field K are equal up to squares if there exists  $c \in K^{\times}$  such that  $x = c^2 y$ .

• We conclude in Section 4 by proving that the local index (given in Section 2) is equal to the index of a conic model (given in Section 3), thereby proving Theorem 1.2.

Acknowledgements. We thank Sergey Finashin and Viatcheslav Kharlamov for helpful correspondence, including explaining the ideas behind Lemma 3.11 to us. We also thank Kirsten Wickelgren for helpful discussions.

Sabrina Pauli acknowledges support by Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) through the Collaborative Research Centre TRR 326 Geometry and Arithmetic of Uniformized Structures, project number 444845124.

Felipe Espreafico acknowledges support of the European Research Council through the grant ROGW-864919.

## 2. The local index

In this section, we will recall the definition and computation of the local index of a line on a degree 2n - 1 hypersurface in  $\mathbb{P}^{n+1}$ . The procedure for computing this index is standard, so the material in this section will be standard as well. It is the interpretation of this index in terms of the geometry at hand that is interesting and will comprise the balance of the article.

Let k be a field. Let  $\mathbb{G}(1, n+1)$  denote the Grassmannian of projective lines in  $\mathbb{P}_k^{n+1}$ . This Grassmannian is isomorphic to G(2, n+2), the Grassmannian of affine 2-planes in affine (n+2)-space. Let  $S \to G(2, n+2)$  denote the tautological bundle, which we can also view as a rank 2 vector bundle  $S \to \mathbb{G}(1, n+1)$ . The rank of the bundle

$$\operatorname{Sym}^{2n-1}(\mathcal{S}^{\vee}) \to \mathbb{G}(1, n+1)$$

is equal to  $\binom{2n-1+2-1}{2n-1} = 2n = \dim \mathbb{G}(1, n+1)$ , so a generic section of this bundle should have a finite vanishing locus. Moreover, any degree 2n - 1 form F on  $\mathbb{P}_k^{n+1}$  determines a section

$$\sigma_F : \mathbb{G}(1, n+1) \to \operatorname{Sym}^{2n-1}(\mathcal{S}^{\vee})$$
$$\ell \mapsto F|_{\ell}$$

that vanishes precisely on those lines contained in the hypersurface  $\mathbb{V}(F)$ . One can then obtain an enriched count of the lines on  $\mathbb{V}(F)$  by computing the Euler number  $e(\operatorname{Sym}^{2n-1}(\mathcal{S}^{\vee})) \in \operatorname{GW}(k)$ , using a Poincaré–Hopf formula to express  $e(\operatorname{Sym}^{2n-1}(\mathcal{S}^{\vee}))$  as a sum of local degrees of  $\sigma_F$  over its vanishing locus, and giving a geometric interpretation to these local indices. The Euler number and Poincaré–Hopf formula for this bundle are given in [BW23, Theorem 1.1 and Corollary 6.9].

In order to compute the local indices of  $\sigma_F$ , we need to make a suitable choice of coordinates on  $\mathbb{G}(1, n + 1)$  and local trivializations of  $\operatorname{Sym}^{2n-1}(\mathcal{S}^{\vee})$ . For coordinates, we give coordinates centered about each line. Given a line  $\ell \in \mathbb{G}(1, n + 1)$ , we base change to the field of definition  $k(\ell)$  and pick coordinates  $[u : v : x_1 : \ldots : x_n]$  of  $\mathbb{P}_{k(\ell)}^{n+1}$  such that  $\ell = \mathbb{V}(x_1, \ldots, x_n)$ . Now [u : v] are the coordinates on  $\ell$ , and we have an open

8

affine  $U_{\ell} \subset \mathbb{G}(1, n+1)$  with coordinates  $(a_1, b_1, \ldots, a_n, b_n)$  corresponding to lines of the (parametric) form

$$\{[u: v: a_1u + b_1v: \ldots: a_nu + b_nv]: [u, v] \in \mathbb{P}^1\}.$$

We obtain a local trivialization of  $\operatorname{Sym}^{2n-1}(\mathcal{S}^{\vee})$  by reading off the coefficients in the monomial basis  $\{u^{2n-1}, u^{2n-2}v, \ldots, v^{2n-1}\}$ .

**Remark 2.1.** Our use of base change to define coordinates may appear problematic, but no problems actually arise. The field of definition  $k(\ell)$  is always a separable extension of k [EH16, Theorem 6.34], and the local index can always be computed by "changing base and taking trace" over separable extensions [BBM+21].

Our chosen coordinates and trivializations need to be compatible (in a precise sense) with a chosen *relative orientation* of the bundle  $\operatorname{Sym}^{2n-1}(\mathcal{S}^{\vee}) \to \mathbb{G}(1, n+1)$ , which is an isomorphism

$$\det \operatorname{Sym}^{2n-1}(\mathcal{S}^{\vee}) \otimes \omega_{\mathbb{G}(1,n+1)} \cong \mathcal{L}^{\otimes 2}$$

for some line bundle  $\mathcal{L} \to \mathbb{G}(1, n+1)$ . Such an isomorphism need not exist for a general vector bundle, but it is a fact that  $\operatorname{Sym}^{2n-1}(\mathcal{S}^{\vee}) \to \mathbb{G}(1, n+1)$  is relatively orientable (see e.g. [OT14, Corollary 7] and [BW23, Corollary 6.9]). We will not explicitly show that our coordinates and trivialization are compatible with this relative orientation, but rather simply refer to [KW21, Section 5] and [Pau22, Section 2.2] for a demonstration of the proof in the n = 2 and n = 3 cases. The n > 3 cases are all completely analogous.

2.1. Computing the local index. Because  $\ell$  is a simple zero of  $\sigma_F$  [EH16, Theorem 6.34], the local index  $\operatorname{ind}_{\ell} \sigma_F$  can be computed by using our coordinates and trivializations to write out  $\sigma_F$  as a polynomial in  $(a_1, b_1, \ldots, a_n, b_n)$  and taking the Jacobian determinant of this polynomial [KW19, Lemma 9] and evaluating at  $(0, 0, \ldots, 0, 0)$ .

By our choice of coordinates  $[u:v:x_1:\ldots:x_n]$  on  $\mathbb{P}^{n+1}$ , we may assume that

$$F = x_1 P_1(u, v) + x_2 P_2(u, v) + \ldots + x_n P_n(u, v) + R(u, v, x_1, \ldots, x_n),$$

where  $P_i$  are homogeneous polynomials of degree 2n - 2 and R is a polynomial in the ideal  $(x_1, \ldots, x_n)^2$ . In our coordinates  $(a_1, b_1, \ldots, a_n, b_n)$  on  $U_\ell$ , the section  $\sigma_F$  becomes the polynomial

$$\sigma_F(\mathbf{a}, \mathbf{b}) = \sum_{i=1}^n (a_i u + b_i v) P_i(u, v) + R(a_1 u + b_1 v, \dots, a_n u + b_n v).$$

The partial derivatives evaluated at  $(0, \ldots, 0)$  are now straightforward to compute:

$$\frac{\partial \sigma_F}{\partial a_i}\Big|_{(0,\dots,0)} = uP_i(u,v) \quad \text{and} \quad \frac{\partial \sigma_F}{\partial b_i}\Big|_{(0,\dots,0)} = vP_i(u,v).$$

To compute the Jacobian matrix, it thus suffices to write out each  $P_i$  in terms of the monomial basis  $\{u^{2n-1}, u^{2n-2}v, \ldots, v^{2n-1}\}$ , as this is our local trivialization of  $\operatorname{Sym}^{2n-1}(\mathcal{S}^{\vee})$ .

So let  $p_{j,i} \in k(\ell)$  be coefficients (for  $1 \le i \le n$  and  $0 \le j \le 2n-2$ ) such that

$$P_i(u,v) = \sum_{j=0}^{2n-2} p_{j,i} u^j v^{2n-2-j}.$$

Then the Jacobian matrix of  $\sigma_F$  at  $(0, \ldots, 0)$  is

$$(2.1) A_{P_1,\dots,P_n} := \begin{pmatrix} p_{2n-2,1} & 0 & p_{2n-2,2} & 0 & \cdots & p_{2n-2,n} & 0 \\ p_{2n-3,1} & p_{2n-2,1} & p_{2n-3,2} & p_{2n-2,2} & \cdots & p_{2n-3,n} & p_{2n-2,n} \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ p_{0,1} & p_{1,1} & p_{0,2} & p_{1,2} & \cdots & p_{0,n} & p_{1,n} \\ 0 & p_{0,1} & 0 & p_{0,2} & \cdots & 0 & p_{0,n} \end{pmatrix}$$

Now

(2.2) 
$$\operatorname{ind}_{\ell} \sigma_F = \operatorname{Tr}_{k(\ell)/k} \langle \det A_{P_1,\dots,P_n} \rangle \in \mathrm{GW}(k).$$

2.2. Examples: lines on cubic surfaces and quintic threefolds. We conclude this section by writing out the Jacobian matrix for the local index in the cases of lines on cubic surfaces and quintic threefolds. We will then sketch how Kass–Wickelgren and Pauli gave geometric interpretations of these determinants and what fails in the general case.

**Example 2.2** (Cubic surfaces). For cubic surfaces, we have  $\operatorname{ind}_{\ell} \sigma_F = \operatorname{Tr}_{k(\ell)/k} \langle \det A_{P_1, P_2} \rangle$  with

$$A_{P_1,P_2} = \begin{pmatrix} p_{2,1} & 0 & p_{2,2} & 0\\ p_{1,1} & p_{2,1} & p_{1,2} & p_{2,2}\\ p_{0,1} & p_{1,1} & p_{0,2} & p_{1,2}\\ 0 & p_{0,1} & 0 & p_{0,2} \end{pmatrix}.$$

Here, we have  $F = x_1P_1 + x_2P_2 + R$  with  $P_i(u, v) = p_{2,i}u^2 + p_{1,i}uv + p_{0,1}v^2$ . Note that det  $A_{P_1,P_2}$  is equal to the resultant of  $P_1$  and  $P_2$ , which is where Kass and Wickelgren's geometric interpretation begins. The Gauß map along  $\ell$  can be identified with the degree 2 map

$$\mathcal{G}: \ell \to \mathbb{P}^1_{k(\ell)}$$
$$[u:v] \mapsto [P_1(u,v): P_2(u,v)].$$

In particular, for each point  $p \in \ell$  there is another point q such that  $T_p \mathbb{V}(F) = T_q \mathbb{V}(F)$ . The Segre involution  $i : \ell \to \ell$  swaps p and q. The fixed points of i are defined over a quadratic field extension  $k(\ell)(\sqrt{\alpha})$  for some  $\alpha \in k(\ell)^{\times}/(k(\ell)^{\times})^2$ , and the Segre index of  $\ell$  is  $\operatorname{Tr}_{k(\ell)/k}\langle \alpha \rangle$ .

To show that this Segre index describes  $\operatorname{ind}_{\ell} \sigma_F$ , we need to show that  $\alpha$  and  $\operatorname{Res}(P_1, P_2)$  agree up to squares. This is proved in [KW21, Proposition 14], but we recall the details here since we use slightly different language. The fixed points of the Segre involution are given by the ramification locus of  $[P_1 : P_2]$ , namely where  $x_1P_2(u, v) - x_2P_1(u, v)$ 

has a multiple root in u and v. The ramification locus is given by the vanishing of the discriminant  $\text{Disc}_{u,v}(x_1P_2(u,v)-x_2P_1(u,v))$ . This locus is defined over  $k(\ell)(\sqrt{\alpha})$ , where

$$\alpha = \operatorname{Disc}_{x_1, x_2}(\operatorname{Disc}_{u, v}(x_1 P_2(u, v) - x_2 P_1(u, v)))$$

We conclude by computing  $\operatorname{Res}(P_1, P_2) = 16 \cdot \alpha$ .

For n > 2, the determinant det  $A_{P_1,\ldots,P_n}$  is not a resultant, which explains why [KW21] does not generalize. However, it turns out that det  $A_{P_1,\ldots,P_n}$  is always a *product* of resultants. We had wondered if this were true after Pauli treated the case of lines on quintic threefolds:

**Example 2.3** (Quintic threefolds). For quintic threefolds, we need to consider the determinant of

$$A_{P_1,P_2,P_3} = \begin{pmatrix} p_{4,1} & 0 & p_{4,2} & 0 & p_{4,3} & 0 \\ p_{3,1} & p_{4,1} & p_{3,2} & p_{4,2} & p_{3,3} & p_{4,3} \\ p_{2,1} & p_{3,1} & p_{2,2} & p_{2,3} & p_{3,3} & p_{3,3} \\ p_{1,1} & p_{2,1} & p_{1,2} & p_{2,2} & p_{1,3} & p_{2,3} \\ p_{0,1} & p_{1,1} & p_{0,2} & p_{1,2} & p_{0,3} & p_{1,3} \\ 0 & p_{0,1} & 0 & p_{0,2} & 0 & p_{0,3} \end{pmatrix}$$

Here, we have  $F = x_1P_1 + x_2P_2 + x_3P_3 + R$  with  $P_i(u, v) = p_{4,i}u^4v + p_{3,i}u^3v^2 + p_{2,i}u^2v^3 + p_{1,i}uv^4 + p_{0,i}v^5$ . Now the image of the Gauß map

$$\mathcal{G}: \ell \to \mathbb{P}^2_{k(\ell)}$$
$$[u:v] \mapsto [P_1(u,v): P_2(u,v): P_3(u,v)]$$

is a rational plane quartic curve. A general rational plane quartic has three nodes  $\nu_1, \nu_2, \nu_3$  defined over the algebraic closure  $\overline{k}$ . Over  $k(\ell)$  we could have 1, 2 or 3 nodes, in each case the sum of the degrees of the residue fields of the nodes over  $k(\ell)$  equals 3. That is,  $\sum_{\text{nodes }\nu} [k(\nu) : k(\ell)] = 3$ .

Each node  $\nu$  defines a degree 2 divisor  $D_{\nu}$  on  $\ell_{k(\nu)}$  as follows. Consider the pencil  $H_t^{\nu}$  of lines in  $\mathbb{P}^2_{k(\nu)}$  through  $\nu$ . Then  $H_t^{\nu} \cap \mathcal{G}(\ell)$  defines a pencil of degree 4 divisors  $D_{\nu} + D_t^{\nu}$  on  $\ell_{k(\nu)}$ .

By [Pau22, Lemma 3.1], the pencil  $D_t^{\nu}$  is base point free when  $\ell$  is a simple (i.e. reduced) line and thus defines a degree 2 map of the form  $[Q_1^{\nu}: Q_2^{\nu}]: \mathbb{P}_{k(\nu)}^1 \to \mathbb{P}_{k(\nu)}^1$ . The non-trivial element of the Galois group of the double cover  $[Q_1^{\nu}: Q_2^{\nu}]$  gives an involution on  $\ell_{k(\nu)}$ , which we again call the *Segre involution*. Geometrically, the Segre involution swaps the pairs of points in the pencil  $D_t^{\nu}$  (see Figure 2A). The fixed points of the Segre involution are defined over  $k(\nu)(\sqrt{\alpha_{\nu}})$  for some  $\alpha_{\nu} \in k(\nu)^{\times}/(k(\nu)^{\times})^2$ , which can be computed as the resultant  $\alpha_{\nu} = \operatorname{Res}(Q_1^{\nu}, Q_2^{\nu})$  by the same argument as above for n = 2. The *index* of this Segre involution is given by the norm

$$\mathcal{N}_{k(\nu)/k(\ell)} \alpha_i = \mathcal{N}_{k(\nu)/k(\ell)} \operatorname{Res}(Q_1^{\nu}, Q_2^{\nu}).$$

We want to identify the local index  $\operatorname{ind}_{\ell}(\sigma_F) = \operatorname{Tr}_{k(\ell)/k} \langle \det A_{P_1,P_2,P_3} \rangle$  with the trace of the product of Segre indices

$$\operatorname{Tr}_{k(\ell)/k} \langle \prod_{\text{nodes } \nu} \mathcal{N}_{k(\nu)/k(\ell)} \operatorname{Res}(Q_1^{\nu}, Q_2^{\nu}) \rangle$$

That is, we need to show that

$$\langle \det A_{P_1,P_2,P_3} \rangle = \langle \prod_{\nu} \mathcal{N}_{k(\nu)/k(\ell)} \operatorname{Res}(Q_1^{\nu}, Q_2^{\nu}) \rangle$$

in  $GW(k(\ell))$ .

In this example we assume that  $\nu_1, \nu_2, \nu_3$  are all defined over  $k(\ell)$ . We can then use a change of coordinates to assume that

$$u_1 = [1:0:0],$$
  
 $u_2 = [0:1:0],$   
 $u_3 = [0:0:1].$ 

Under this assumption, the quartic polynomials  $P_i, P_j$  have two common zeros for  $i \neq j$ , and hence there exist quadratic polynomials  $Q_1, Q_2, Q_3$  such that

$$P_1 = Q_2 Q_3,$$
  
 $P_2 = Q_1 Q_3,$   
 $P_3 = Q_1 Q_2.$ 

In this case, one can prove that

$$\begin{split} & [Q_1^1:Q_2^1] = [Q_2:Q_3], \\ & [Q_1^2:Q_2^2] = [Q_1:Q_3], \\ & [Q_1^3:Q_2^3] = [Q_1:Q_2] \end{split}$$

and det  $A_{P_1,P_2,P_3} = \prod_{i < j} \text{Res}(Q_i, Q_j)$ , thereby proving that the local index is the product of the Segre indices.

Our use of coordinate change to write  $\nu_1, \nu_2, \nu_3$  in this simple form relies on the assumption that these nodes are all defined over  $k(\ell)$ . This need not be the case — in general,  $\nu_1, \nu_2, \nu_3$  will be Galois conjugates over  $\overline{k}$ . We can then use a coordinate change to again express  $P_1, P_2, P_3$  as products of quadratic polynomials  $Q_1, Q_2, Q_3$ , with these quadratics being Galois conjugate. See [Pau22] for more details.

For n > 3, we can no longer use a projective change of coordinates to reduce to a particular case as was done for quintic threefolds. This is the reason that [Pau22] does not admit an obvious generalization to lines on hypersurfaces of greater dimension and degree. A construction of Finashin and Kharlamov will allow us to sidestep this issue. This construction uses what we call *conic models*, which we introduce in the next section.

#### QUADRATIC SEGRE INDICES

## 3. Conic models for the Gauss curve

Our goal in this section is to define *conic models* for Gauß curves. This construction appears in [FK21, §6], and we will use it to show that the local index is the same as the product of Segre indices. In particular, this construction allows us to see that the determinant det  $A_{P_1,\dots,P_n}$  (see Equation 2.1) is always a product of resultants.

**Remark 3.1.** For the remainder of the article, we will assume  $n \ge 3$ . Most of the constructions from here on out only make sense under this assumption.

We start by recalling the definition of the Segre index in general, following the same notation used in Examples 2.2 and 2.3. This was already defined in the real case in [FK21] and there are no major technical differences in passing to a general field. The main tool we will need is a result attributed to Castelnuovo, which states that there are finitely many (n-3)-dimensional (2n-4)-secants to a generic degree 2n-2 rational curve in  $\mathbb{P}^{n-1}$ . This generalizes the fact that a general rational quartic plane curve has three nodes. Segre involutions and their associated indices are constructed from these finite sets of secants.

Finally, we will dive into conic models for Gauß curves: to each Gauß curve and its Castelnuovo secants, we can associate a set of points B and a conic Q in  $\mathbb{P}^2_{k(\ell)}$ . We will show that the local index and the Segre index can be computed purely in terms of a conic model.

3.1. Secants and the Segre index. Recall our setting. Let  $X \subset \mathbb{P}^{n+1}$  be a generic hypersurface of degree 2n - 1, and fix a line  $\ell \subseteq X$ . Without loss of generality, we may choose coordinates  $[u : v : x_1 : \ldots : x_n]$  on  $\mathbb{P}^{n+1}_{k(\ell)}$  such that  $\ell = \{[u : v : x_1 : \ldots : x_n] : x_1 = \cdots = x_n = 0\}$  and  $X = \mathbb{V}(F)$  for

$$F = x_1 P_1(u, v) + x_2 P_2(u, v) + \ldots + x_n P_n(u, v) + R(u, v, x_1, \ldots, x_n),$$

where  $P_1, \ldots, P_n \in k(\ell)[u, v]$  are homogeneous polynomials of degree 2n - 2 and  $R \in (x_1, \ldots, x_n)^2 \subseteq k(\ell)[u, v, x_1, \ldots, x_n]$  is also of degree 2n - 2.

Following [FK21], the Gauß map along  $\ell$  is given by

(3.1) 
$$\mathcal{G}: \mathbb{P}^{1}_{k(\ell)} \to \mathbb{P}^{n-1}_{k(\ell)}$$
$$[u:v] \mapsto [P_{1}(u,v):\ldots:P_{n}(u,v)].$$

**Definition 3.2.** We call the image of  $\ell$  under the Gauß map the *Gauß curve* associated to  $\ell$ . This is a rational curve

 $\mathcal{G}(\ell) \subseteq \mathbb{P}^{n-1}$ 

of degree 2n-2.

We are interested in the (n-3)-dimensional (2n-4)-secants to the Gauß curve, i.e. the (n-3)-planes that meet the Gauß curve in 2n-4 points (when counted with multiplicity). Finashin–Kharlamov state that there are (geometrically)  $\binom{n}{2}$  such secants [FK21, Proposition 4.3.3], which they refer to as the *Castelnuovo count*.

**Proposition 3.3** (Castelnuovo count). The number of (n-3)-dimensional (2n-4)-secants (defined over  $\overline{k}$ ) to a generic rational curve  $C \subset \mathbb{P}^{n-1}$  of degree 2n-2 is  $\binom{n}{2}$ .

*Proof.* Just as described in the proof of [FK21, Proposition 4.3.3 (2)], the number of secants (when counted with multiplicity) is  $\binom{n}{2}$ , and these multiplicities are always positive. This fact follows from the same argument present in [EH16, §12.4.4], where Eisenbud and Harris apply Porteous' formula to count the number of quadrisecant lines to rational curves in  $\mathbb{P}^3$  (i.e. the case n = 4). We explain how this argument goes in general.

Recall that the parameter space for dimension 0, degree 2n - 4 subschemes  $\Gamma \subseteq \mathbb{P}^1$  is given by  $\operatorname{Sym}^{2n-4} \mathbb{P}^1 \cong \mathbb{P}^{2n-4}$ . Consider the vector bundle  $\mathcal{E}^* \to \mathbb{P}^{2n-4}$  whose fibers are given by

$$\mathcal{E}_{\Gamma}^* = H^0(\mathcal{O}_{\mathbb{P}^1}(2n-2))/H^0(\mathcal{I}_{\Gamma}(2n-2)),$$

which is the space polynomials of degree 2n - 2 modulo the polynomials that vanish on  $\Gamma$ . Note that the rank of  $\mathcal{E}^*$  is 2n - 1 - 3 = 2n - 4.

The Gauß map  $\mathcal{G}: \ell \to \mathbb{P}^{n-1}$  induces a map

$$H^0(\mathcal{O}_{\mathbb{P}^{n-1}}(1)) \hookrightarrow H^0(\mathcal{O}_{\mathbb{P}^1}(2n-2)),$$

given by simply sending  $x_i$  to  $P_i$  (in the notation defined at the beginning of this subsection). Denoting by  $\mathcal{F}$  the trivial rank *n* bundle over  $\mathbb{P}^{2n-4}$  with fiber  $H^0(\mathcal{O}_{\mathbb{P}^{n-1}}(1))$ , we get a morphism of vector bundles

$$\varphi: \mathcal{F} \to \mathcal{E}^*$$

by projecting to the quotient. The (2n-4)-secants to C of dimension n-3 correspond to the locus of dimension 0, degree 2n-4 subschemes  $\Gamma \subseteq C \subseteq \mathbb{P}^{n-1}$  whose projective linear span has codimension 2. Given  $\Gamma \subseteq C \subseteq \mathbb{P}^{n-1}$ , by the definition of the  $\varphi$ , the equations in the kernel of  $\varphi_{\Gamma}$  correspond to hyperplanes that contain  $\Gamma$  and, therefore, the linear subspace determined by the equations in ker  $\varphi$  is the projective span of  $\Gamma$ . To have a projective span of codimension two, we need ker  $\varphi$  to have dimension 2 and therefore rank  $\varphi = n - 2$ . It thus suffices to consider the locus  $M_{n-2}(\varphi) \subseteq \mathbb{P}^{2n-4}$  where the map has rank n-2.

The expected codimension of  $M_m(\varphi)$  is (e-m)(f-m), where e and f are the ranks of  $\mathcal{E}^*$  and  $\mathcal{F}$  respectively. This gives us that the expected dimension of  $M_m(\varphi)$  is 2n - (2n - 4 - m)(n - m). Notice that plugging in m = n - 2 gives us dimension zero, which means that generic rational curves of degree 2n - 2 in  $\mathbb{P}^{n-1}$  have a finite number of (n - 3)-dimensional (2n - 4)-secants. From this formula, one can also verify that the locus, where  $\varphi$  has rank less than n - 2, has expected negative dimension.

It remains to calculate the class  $[M_{n-2}(\varphi)]$  in the Chow ring of  $\mathbb{P}^{2n-4}$ . This is done by Porteous' formula [EH16, Theorem 12.4]. In the present context, Porteous' formula and

the computation of the total Chern class  $c(\mathcal{E}^*)$  [EH16, Theorem 10.16] give us

$$[M_{n-2}(\varphi)] = \begin{vmatrix} c_{n-2}(\mathcal{E}^*) & c_{n-1}(\mathcal{E}^*) \\ c_{n-3}(\mathcal{E}^*) & c_{n-2}(\mathcal{E}^*) \end{vmatrix}$$
$$= \begin{vmatrix} \binom{n}{2}\zeta^{n-2} & \binom{n+1}{2}\zeta^{n-1} \\ \binom{n-1}{2}\zeta^{n-3} & \binom{n}{2}\zeta^{n-2} \end{vmatrix}$$
$$= \left( \binom{n}{2}^2 - \binom{n+1}{2}\binom{n-1}{2} \right) \zeta^{2n-4},$$

where  $\zeta$  is the hyperplane class. We now conclude by noting that  $\binom{n}{2}^2 - \binom{n+1}{2}\binom{n-1}{2} = \binom{n}{2}$ .

For our purposes, we also need each of these secants to have multiplicity 1 for a generic C. Again, the proof proceeds exactly as for [FK21, Proposition 4.3.3 (3)] — one uses Kleiman's transversality theorem [Kle74] to show that over  $\overline{k}$ , there is a dense open subset of the space of degree 2n - 2 rational curves in  $\mathbb{P}^{n-1}$  on which all secants have multiplicity 1, since these multiplicities arise as an intersection multiplicity in an appropriate parameter space.

**Example 3.4.** When n = 3, we recover the three nodes on a rational plane quartic as secants (see Figure 2A). Indeed, we get three 2-secants of dimension zero. These are points that intersect the curve twice, i.e. are double points. When n = 4, we get six 4-secants of dimension 1 to a rational sextic in  $\mathbb{P}^3$  (see Figure 2B).

We now define the Segre involution associated to a secant of the Gauß curve. Let S be one of the (2n - 4)-secants to  $\mathcal{G}(\ell)$ , and let  $k(S)/k(\ell)$  be its field of definition. Then Sdefines a degree 2n - 4 divisor  $D_S$  on  $\ell_{k(S)}$ , as  $S \cap \mathcal{G}(\ell)$  consists of 2n - 4 geometric points (when counted with multiplicity). Now consider the pencil  $H_t$  of hyperplanes in  $\mathbb{P}_{k(S)}^{n-1}$ containing S. For each t, we get a degree 2n - 2 divisor on  $\ell_{k(S)}$ , as  $H_t \cap \mathcal{G}(\ell)$  consists of 2n - 2 geometric points (when counted with multiplicity) by Bézout's theorem. We can write this divisor as  $D_S + D_t^S$ . Note that  $D_t^S$  is base point free and therefore defines a degree 2 map  $[Q_1^S : Q_2^S] : \mathbb{P}_{k(S)}^1 \to \mathbb{P}_{k(S)}^1$ .

**Definition 3.5.** We define the Segre involution associated to a secant S of  $\mathcal{G}(\ell)$  as the involution

$$i_S: \ell_{k(S)} \to \ell_{k(S)}$$

given by the non-trivial element of the Galois group of the double covering  $[Q_1^S:Q_2^S]$ .

The fixed points of the Segre involution  $i_S$  are defined over  $k(S)(\sqrt{\alpha_S})$  for some  $\alpha_S \in k(S)^{\times}/(k(S)^{\times})^2$ . In fact, we may take  $\alpha_S = \text{Res}(Q_1^S, Q_2^S)$  by the argument given in Example 2.2.

**Definition 3.6.** The Segre index of  $\ell$  is

$$\operatorname{seg}(X, \ell) := \operatorname{Tr}_{k(\ell)/k} \langle \prod_{S} \operatorname{N}_{k(S)/k(\ell)} \alpha_{S} \rangle \in \operatorname{GW}(k)$$

where the product goes over all the secants S. As in Example 2.3, each secant S is a representative from its orbit of Galois conjugates whose contributions to  $seg(X, \ell)$  are encoded in the norm  $N_{k(S)/k(\ell)} \alpha_S$ .

**Example 3.7.** In case  $k = k(\ell) = \mathbb{R}$  the Segre index  $\operatorname{seg}(X, \ell)$  agrees with the Segre index from the introduction defined by Finashishin–Kharlamov (you just have to stick brackets around it). Indeed, each secant defined over  $\mathbb{C}$  will contribute the factor  $N_{\mathbb{C}/\mathbb{R}}(1) = 1$  and each secant S defined over  $\mathbb{R}$  will contribute the factor  $\alpha_S$ .

**Remark 3.8.** As mentioned in the introduction, we will generally conflate the Segre involution  $i_S : \ell_{k(S)} \to \ell_{k(S)}$ , which is an involution of a projective line  $\mathbb{P}^1_{k(S)}$ , with an involution of the Gauß curve  $\mathcal{G}(\ell)_{k(S)}$ . In order to justify this conflation, we need to verify that  $i_S$  induces an involution on  $\mathcal{G}(\ell)_{k(S)}$  whose fixed points are defined over  $k(S)(\sqrt{\alpha_S})$ . To define an involution on  $\mathcal{G}(\ell)_{k(S)}$ , we first note that  $\mathcal{G}(\ell)_{k(S)} = \mathcal{G}(\ell_{k(S)})$ . We may therefore define

$$i'_{S}: \mathcal{G}(\ell)_{k(S)} \to \mathcal{G}(\ell)_{k(S)}$$
$$\mathcal{G}(x) \mapsto \mathcal{G}(i_{S}(x)).$$

That is,  $i'_S = \mathcal{G} \circ i_S$ . Note that  $i_S(x) = x$  only if x and  $i_S(x)$  have the same image under the Gauß map — this is by definition of the involution  $i_S$ . It follows that the fixed points of  $i'_S$  are given by  $\{\mathcal{G}(x) : i_S(x) = x\}$ . Finally, since  $\mathcal{G} : \ell_{k(S)} \to \mathcal{G}(\ell)_{k(S)}$  is a birational map, we have an isomorphism from the field of definition of the fixed locus of  $i_S$  to the field of definition of the fixed locus of  $i'_S$ , as desired.

3.2. Conic models. Our next goal is to recap Finashin and Kharlamov's construction of *conic models* for the Gauß curve [FK21, Section 6.3]. These consist of a plane conic Q, together with a collection a zero dimensional subscheme  $B \subseteq \mathbb{P}^2$  of degree  $\binom{n}{2}$ , such that the Gauß curve coincides with the strict transform of Q in the blowup  $\operatorname{Bl}_B \mathbb{P}^2$ .

One place where we deviate from Finashin–Kharlamov is that we require our plane conic Q to come equipped with a parameterization, which we need in order to define an index associated to the conic. Such a parameterization comes from stereographic projection from a k-rational point on Q, but over non-closed fields there are conics defined over k with no k-points. Working with the space of *parameterized conics* allows us to sidestep this issue.

**Remark 3.9.** Technically, Finashin and Kharlamov implicitly assume that their conics are parameterized, as they require their conics to have real points in order to describe the associated conic index. However, their comment on this construction is brief enough that they do not justify why one can always take a parameterized conic model. We will give such a justification in the course of this section.

Let B be a zero dimensional subscheme of  $\mathbb{P}^2$  of degree  $\binom{n}{2}$  of  $\binom{n}{2}$  (geometric) points in  $\mathbb{P}^2$  in general position, and let  $Q \subseteq \mathbb{P}^2$  be a conic that does not pass through any of the

points in B. Curves of degree n-1 through B form a linear system  $L_B$  of dimension

$$\dim L_B = \binom{3+n-1-1}{n-1} - \binom{n}{2} - 1$$
$$= \binom{n+1}{2} - \binom{n}{2} - 1$$
$$= n-1.$$

Therefore, after choosing a basis  $\beta$  for this system, we obtain a rational map

$$g_{B,\beta}: \mathbb{P}^2 \dashrightarrow \mathbb{P}^{n-1}$$

whose indeterminacy locus is B. Of course, such a map depends on the choice of a basis. Finally, the curve  $C := g_{B,\beta}(Q) \subset \mathbb{P}^{n-1}$  is a rational curve of degree 2n - 2.

We can summarize the construction  $g_{B,\beta}(Q)$  in terms of a rational map to the space of rational curves of degree 2n - 2. Let  $\operatorname{Mor}(\mathbb{P}^1, \mathbb{P}^N)_d$  denote the space of rational curves of degree d in  $\mathbb{P}^N$ . Let  $\mathcal{Q} := \operatorname{Mor}(\mathbb{P}^1, \mathbb{P}^2)_2$  denote the space of parameterized conics in  $\mathbb{P}^2$ , and let  $\operatorname{Conf}_m(\mathbb{P}^2)$  denote the configuration space of m points in  $\mathbb{P}^2$ . The space of bases for a linear system of dimension n - 1 is a projective space of dimension  $n^2 - 1$ . Altogether, Finashin–Kharlamov's construction is a rational map of the form

FK : 
$$\operatorname{Conf}_{\binom{n}{2}}(\mathbb{P}^2) \times \mathbb{P}^{n^2-1} \times \mathcal{Q} \dashrightarrow \operatorname{Mor}(\mathbb{P}^1, \mathbb{P}^{n-1})_{2n-2}$$
  
 $(B, \beta, Q) \mapsto g_{B,\beta}(Q).$ 

**Definition 3.10.** A conic model for a rational curve  $C \in Mor(\mathbb{P}^1, \mathbb{P}^{n-1})_{2n-2}$  is an element of the fiber  $FK^{-1}(C)$ . We say that C has a conic model if the fiber  $FK^{-1}(C)$  is not empty.

As described in [FK21, p. 4077], there is a close relationship between a rational curve with its secants and a conic model for the curve. We now outline this relationship, which is summarized in Table 1.

As previously described, our Gauß curve  $\mathcal{G}(\ell)$  is the image of the plane conic Q in a chosen conic model  $(B, \beta, Q)$ . (The fact that a generic Gauß curve has a conic model will be proved in Lemma 3.11.) Let  $b \in B(\overline{k})$  with residue field k(b). By Cramer's theorem on algebraic curves, there exists a unique plane curve  $Z_b$  of degree n-2 passing through all points of  $B(\overline{k}) - \{b\}$ . The field of definition of  $Z_b$  is k(b). Indeed,  $B(\overline{k}) - \{b\}$  is defined over k(b), since both b and B are defined over k(b). Thus  $B(\overline{k}) - \{b\}$  and hence  $Z_b$  are fixed under Gal(k(b))-action, giving us that  $Z_b$  is defined over k(b).

Bézout's theorem implies that  $Z_b$  and Q intersect in 2n - 4 points. Consequently, the projective linear span of  $g_{B,\beta}(Z_b)$  forms an (n-3)-dimensional (2n-4)-secant to  $\mathcal{G}(\ell)$ . By verifying that distinct points  $b, b' \in B(\overline{k})$  correspond to distinct (n-3)-planes, we conclude that all  $\binom{n}{2}$  such secants to  $\mathcal{G}(\ell)$  arise in this manner.

In particular, this construction establishes a bijection between  $B(\overline{k})$  and the set of (n-3)dimensional geometric (2n-4)-secants to  $\mathcal{G}(\ell)$ . This bijection maps Galois conjugate

points in  $B(\overline{k})$  to Galois conjugate secants, while preserving the associated Galois action. This implies that a point  $b \in B$  corresponds to a secant S satisfying  $k(b) \cong k(S)$ .

Next, take the pencil  $H_t^b$  of lines in  $\mathbb{P}^2$  through a fixed  $b \in B$ . Then  $H_t^b \cup Z_b$  is a pencil of degree n-1 curves through B, each of which consists of a fixed component  $Z_b$  and a moving linear component. The projective linear span of  $g_{B,\beta}(H_t^b \cup Z_b)$  is a pencil of (n-2)-dimensional hyperplanes in  $\mathbb{P}^{n-1}$ , each of which contains the (2n-4)-secant corresponding to b. This pencil of (n-2)-dimensional hyperplanes is used to construct the Segre involution

$$i_S: \mathcal{G}(\ell) \to \mathcal{G}(\ell).$$

Pulling back  $i_S$  under  $g_{B,\beta}$  gives an involution

 $(3.2) \qquad \qquad \mu_b: Q \to Q.$ 

Every involution on a conic can be obtained by intersecting the conic with a pencil of lines through some point, known as the *polar point*, and swapping the pairs of points for each line. One can check that  $\mu_b$  is the involution given by the pencil of lines through b, so that b is the polar point of this involution.

Gauß curves	Conic models
$\mathcal{G}(\ell)$	Q
Secant $S$	$b \in B$
Hyperplanes containing $S$	Lines through $b$
Segre involution $i_S$	Conic involution $\mu_b$

TABLE 1. Dictionary for Gauß curves and their conic models

We will now show that a generic rational curve of degree 2n - 2 in  $\mathbb{P}^{n-1}$  has a conic model, as claimed in [FK21, Section 6.3].

**Lemma 3.11.** Let C be a general rational curve of degree 2n - 2 in  $\mathbb{P}^{n-1}$ . Then there exists  $(B, \beta, Q) \in \operatorname{Conf}_{\binom{n}{2}}(\mathbb{P}^2) \times \mathbb{P}^{n^2-1} \times \mathcal{Q}$  such that  $g_{B,\beta}(Q) = C$ .

Proof. The space  $\operatorname{Mor}(\mathbb{P}^1, \mathbb{P}^N)_d$  is given by  $U/\operatorname{PGL}_2$ , where  $U \subset \mathbb{P}(H^0(\mathbb{P}^1, \mathcal{O}(d))^{\oplus N+1})$ . One can prove that  $\operatorname{Mor}(\mathbb{P}^1, \mathbb{P}^N)_d$  is an irreducible scheme of dimension (d+1)(N+1)-4. To compute this dimension, note that the space of homogeneous degree d polynomials in 2 variables is d+1, and we need N+1 such polynomials to define a morphism to  $\mathbb{P}^N$ . We then subtract 1 to account for projectivization, and we subtract 3 to account for the action of  $\operatorname{Aut}(\mathbb{P}^1) \cong \operatorname{PGL}_2$ . So for N = n - 1 and d = 2n - 2, the space  $\operatorname{Mor}(\mathbb{P}^1, \mathbb{P}^N)_d$ has dimension  $2n^2 - n - 4$ . Note that dim  $\mathcal{Q} = 5$  and dim  $\operatorname{Conf}_{\binom{n}{2}}(\mathbb{P}^2) = 2\binom{n}{2} = n^2 - n$ .

The source and target of FK are both geometrically irreducible (as products of geometrically irreducible schemes). As previously computed, the source has dimension  $(n^2 - n) + 5 + (n^2 - 1) = 2n^2 - n + 4$ , while the target has dimension  $2n^2 - n - 4$ . To prove the desired claim, it remains to show that generic fibers of FK have dimension

$$(2n^{2} - n + 4) - (2n^{2} - n - 4) = 8.$$

To this end, we will show that  $g_{B_1,\beta_1}(Q_1) = g_{B_2,M\beta_2}(Q_2)$  for some change of basis matrix M if and only if there exists a projective transformation of  $\mathbb{P}^2$  transforming  $(B_1, Q_1)$  to  $(B_2, Q_2)$ . From this, it will follow that generic fibers of FK have dimension dim  $\operatorname{Aut}(\mathbb{P}^2) = \operatorname{dim} \operatorname{PGL}_3 = 8$ , as desired.

To begin, assume that  $(B_1, Q_1)$  and  $(B_2, Q_2)$  are projectively equivalent. The curves  $g_{B_j,\beta_j}(Q_j)$  are obtained by embedding the strict transform of  $Q_j$  on the blowup  $\operatorname{Bl}_{B_j}(\mathbb{P}^2)$ , and an automorphism taking  $(B_1, Q_1)$  to  $(B_2, Q_2)$  gives an isomorphism of the strict transforms of  $Q_j$  on  $\operatorname{Bl}_{B_j}(Q_j)$ . Now the embeddings  $g_{B_j,\beta_j}(Q_j)$  are isomorphic but need not be equal (as elements of  $\operatorname{Mor}(\mathbb{P}^1, \mathbb{P}^{n-1})_{2n-1}$ ), but they will differ by a projective change of coordinates on  $\mathbb{P}^{n-1}$ . Let  $M \in \operatorname{PGL}_n$  represent this change of coordinates. Then  $g_{B_1,\beta_1}(Q_1) = g_{B_2,M\beta_2}(Q_2)$ , as desired.

We now explain why  $g_{B_1,\beta_1}(Q_1) = g_{B_2,\beta_2}(Q_2)$  implies that  $(B_1,Q_1)$  and  $(B_2,Q_2)$  are projectively equivalent. As previously described, there is a natural bijection between  $B_j$  and the set of (n-3)-dimensional (2n-4)-secants to the curve  $g_{B_j,\beta_j}(Q_j)$ . Each Segre involution  $i_S$  on this curve determines an involution  $\mu_b : Q_j \to Q_j$  with polar point  $b \in B_j$ . In particular, we can recover the set  $B_j$  from the curve  $g_{B_j,\beta_j}(Q_j)$  via the involutions  $i_S$  — assuming we already know  $Q_j$ . The polar point of an involution on  $Q_j$ is determined up to projective change of coordinates, so the ambiguity in reconstructing  $(B_j, Q_j)$  from  $g_{B_j,\beta_j}(Q_j)$  is precisely  $\operatorname{Aut}(\mathbb{P}^2) \cong \operatorname{PGL}_3$ . In other words, if  $g_{B_1,\beta_1}(Q_1) =$  $g_{B_2,\beta_2}(Q_2)$ , then there exists a projective transformation taking  $(B_1, Q_1)$  to  $(B_2, Q_2)$ , as desired.

**Remark 3.12.** Lemma 3.11 states that a general rational curve of degree 2n-2 in  $\mathbb{P}^{n-1}$  has a conic model. For n = 4, this description is explicitly related to Dye's result on sextic space curves and double sixes of cubic surfaces [Dye97], as mentioned in Footnote 1. Indeed, blowing up  $\binom{4}{2}$  points in general position in  $\mathbb{P}^2$  yields a cubic surface. The strict transforms of the six conics through five of these points form one half of a double six. Now take any conic in  $\mathbb{P}^2$  that does not meet any of the six points at the center of our blowup. The strict transform of this conic is a sextic curve on the cubic surface, and it meets each of the aforementioned lines in four points. Dye proves that every smooth rational space sextic and its six quadrisecants can be constructed in this manner.

Using the above, one can explicitly construct rational space sextics and their quadrisecants. All that remains is to give a parameterization

$$\mathbb{P}^2 \dashrightarrow \mathbb{P}^3$$

of the cubic surface obtained by blowing up six points. The classical method for constructing such a parameterization is to pick a basis  $f_1, f_2, f_3, f_4$  of the space of cubic forms interpolating the six points in  $\mathbb{P}^2$ . The resulting cubic surface is then the Zariski closure of the map

$$[f_1:\ldots:f_4]:\mathbb{P}^2\dashrightarrow\mathbb{P}^3.$$

For example, the sextic in Figure 2B is the image of the conic  $\mathbb{V}(\frac{x^2}{4} + \frac{y^2}{9} - z^2)$  on a Clebsch cubic surface, which is obtained by blowing up the points

$$B = \{ [1:0:0], [0:1:0], [0:0:1], [1:1:1], [1:2:3], [2:-1:1] \} \}$$

For our basis of cubic forms through these points, we chose

$$f_{1} = \frac{1}{7}(3x^{2}z + xz^{2} + x^{2}y - 5xy^{2}),$$
  

$$f_{2} = \frac{1}{5}(2x^{2}z + xyz - 3xy^{2}),$$
  

$$f_{3} = \frac{1}{2}(-5xz^{2} + 2yz^{2} + 3x^{2}z),$$
  

$$f_{4} = \frac{1}{2}(x^{2}z - 3xz^{2} + 2y^{2}z),$$

which were adapted from [CSD07, Example 7]. The quadrisecant depicted in Figure 2B is the image of the conic through  $B - \{[1:2:3]\}$ .

Not only does every general rational curve admit a conic model, but we can further take our conic and locus of points to have the same field of definition as the rational curve.

**Corollary 3.13.** Let C be a general rational curve of degree 2n - 2 in  $\mathbb{P}^{n-1}$ . If C is defined over a field K/k, then there exists a conic model  $(B, \beta, Q)$  of C such that B and Q are defined over K.

*Proof.* Let  $q : \mathbb{P}^1 \to \mathbb{P}^2$  be any parameterized conic defined over K, and denote  $Q := q(\mathbb{P}^1)$ . We will show that there exist  $(B, \beta) \in \operatorname{Conf}_{\binom{n}{2}}(\mathbb{P}^2) \times \mathbb{P}^{n^2-1}$  (with B defined over K) such that  $g_{B,\beta}(Q) = C$ , which will give the desired claim.

Each secant S to C determines a Segre involution  $i_S : \mathbb{P}^1_{k(S)} \to \mathbb{P}^1_{k(S)}$ , so  $q \circ i_S$  is an involution of  $Q_{k(S)}$ . Let b(S) denote the polar point of  $q \circ i_S$ . Let B denote the scheme whose underlying set is  $\{b(S) : S \text{ secant to } C\}$ . Note that B is defined over K. Indeed, all of the schemes involved are geometrically reduced closed subschemes of projective space, so they are defined over K if and only if they are fixed under  $\operatorname{Gal}(K^{\operatorname{sep}}/K)$ -action (see e.g. [McK25, Proposition 2.4]). The curve C is defined over K, so it and its scheme of secants are fixed under all  $\operatorname{Gal}(K^{\operatorname{sep}}/K)$ -actions. Since we have assumed that Q is defined over K, this conic is also fixed under  $\operatorname{Gal}(K^{\operatorname{sep}}/K)$ -actions, so the set of polar points of the form b(S) must also be  $\operatorname{Gal}(K^{\operatorname{sep}}/K)$ -fixed.

It remains to show that there exists  $\beta$  such that  $g_{B,\beta}(Q) = C$ . Let  $(B', \beta', Q')$  be a conic model for C. Since Q' and C are geometrically birational, they are birational after some extension K' of K. Thus K' is a field of definition of Q'. Since Q' is a degree 2 curve and the characteristic of K is not 2, we know that K'/K is a separable extension. We therefore have a projective transformation M (over K') such that MQ' = Q. The set of involutions induced by MB' and B must be the same, as these are both induced by the involutions  $i_S$  coming from C and its secants. In particular, the polar points of the involutions induced by MB' and B must agree, so MB' = B. It follows that  $(B, M\beta', Q)$ is a conic model for C.

**Definition 3.14.** Given a line  $\ell \subseteq X$ , we will say that a conic model  $(B, \beta, Q)$  of  $\mathcal{G}(\ell)$  is *rational* if both B and Q are defined over  $k(\ell)$ . Corollary 3.13 states that a rational conic model always exists.

3.3. Two equivalent formulas for the Segre index. We have already defined the Segre index of a line  $\ell$  in terms of the fixed points of the Segre involutions associated to the secants to the Gauß curve  $\mathcal{G}(\ell)$ . Using the dictionary between the Gauß curve and its conic models, we can define an alternative index associated to  $\ell$ .

**Definition 3.15.** Let  $(B, \beta, Q)$  be a rational conic model for a Gauß curve  $\mathcal{G}(\ell)$ . For each  $b \in B$ , let  $k(b) \supset k(\ell)$  be the field of definition of  $b \in B$  and let  $\alpha_b \in k(b)^{\times}/(k(b)^{\times})^2$ be such that the fixed points of the involution  $\mu_b : Q \to Q$  (see Equation 3.2) are defined over  $k(\sqrt{\alpha_b})$ . The *conic index* of  $\ell$  is

$$\operatorname{con}(X,\ell) := \operatorname{Tr}_{k(\ell)/k} \langle \prod_{b \in B} \operatorname{N}_{k(b)/k(\ell)} \alpha_b \rangle \in \operatorname{GW}(k).$$

Note that the conic index does not depend on our choice of conic model for  $\mathcal{G}(\ell)$ . Indeed, any two such choices differ by a projective change of coordinates over  $k(\ell)$  (as described in the proof of Lemma 3.11 and Corollary 3.13), and such a change of coordinates does not change the field of definition of the fixed points of an involution of the conic.

We will conclude this section by showing that the Segre index and the conic index are equal.

**Proposition 3.16.** Given a line  $\ell$  on a degree 2n - 1 hypersurface  $X \subseteq \mathbb{P}^{n+1}$ , we have  $seg(X, \ell) = con(X, \ell)$ .

Proof. Fix a rational conic model  $(B, \beta, Q)$  for  $\mathcal{G}(\ell)$ . We have seen that if S is the secant corresponding to  $b \in B$ , then  $g_{B,\beta}$  induces a field isomorphism  $\phi : k(S) \to k(b)$  that fixes  $k(\ell)$ . Furthermore, we have  $\phi(\alpha_S) = \alpha_b$  up to squares simply because the involution in the conic model is mapped to the involution of the Gauß curve by  $g_{B,\beta}$ .

**Example 3.17** (The conic index for lines on a quintic threefold). We look at the case n = 3 (the quintic threefold case). In this case the map  $g_{B,\beta} \colon \mathbb{P}^2 \dashrightarrow \mathbb{P}^{n-1} = \mathbb{P}^2$  is a birational map and therefore a Cremona transformation. Recall that the Gauß map  $\mathcal{G} \colon \mathbb{P}^1_{k(\ell)} \to \mathbb{P}^2_{k(\ell)}$  in this case is a general degree 4 parametrized plane curve and that it has three nodes. These nodes are exactly the  $\binom{n}{2} = 3$  zero dimensional 2n - 4-secants. Assume for simplicity that the nodes are all defined over  $k(\ell)$  and lie in general position (just like we did in Example 2.3). Then, after a coordinate change we can assume that the nodes are the points [1:0:0], [0:1:0] and [0:0:1]. The Cremona transformation) is given by  $\operatorname{Cr} := [x_2x_3: x_1x_3: x_1x_2] \colon \mathbb{P}^2 \dashrightarrow \mathbb{P}^2$ . Now  $\operatorname{Cr} \circ \mathcal{G} : \mathbb{P}^1_{k(\ell)} \to \mathbb{P}^2_{k(\ell)}$  has degree 2, so we have a parametrized conic  $Q := \operatorname{Cr} \circ \mathcal{G}$ . Let  $B = \{[1:0:0], [0:1:0], [0:0:1]\}$  and  $g_{B,\beta} = \operatorname{Cr}$ . Then  $(B, \beta, Q)$  is a conic model for  $\mathcal{G}$  since  $\operatorname{Cr}$  is a birational involution with base locus B.

We compute the conic index in this setup. To do so, we first have to write down a parametrization of  $Q = \operatorname{Cr} \circ \mathcal{G}$ . Recall from Example 2.3 that in this case, the Gauß map  $\mathcal{G}$  is given by  $\mathcal{G} = [Q_2Q_3 : Q_1Q_3 : Q_1Q_2]$  and thus  $\operatorname{Cr} \circ \mathcal{G} = [Q_1 : Q_2 : Q_3]$ . For

the polar point  $[1:0:0] \in B$  the pencil of degree two divisors on  $\mathbb{P}^1_{k(\ell)}$  is given by  $\{t_0Q_2+t_1Q_3=0\} \subset \mathbb{P}^1_{k(\ell)}$  and thus the fixed points of the associated involution live over  $k(\ell)(\sqrt{\alpha})$  with  $\alpha = \operatorname{Res}(Q_2, Q_3)$  by the same argument as in Example 2.2. Similarly, the fixed points of the involutions for the other two points in B live over  $k(\ell)(\sqrt{\alpha})$  with  $\alpha = \operatorname{Res}(Q_1, Q_3)$  respectively  $\alpha = \operatorname{Res}(Q_1, Q_2)$ . Thus the conic index equals

$$\operatorname{Res}(Q_2, Q_3) \operatorname{Res}(Q_1, Q_3) \operatorname{Res}(Q_1, Q_2),$$

which agrees with the Segre index and local index by Example 2.3.

## 4. The local index is the conic index

We are almost ready to prove Theorem 1.2, which states that the local index of a line on a degree 2n - 1 hypersurface in  $\mathbb{P}^{n+1}$  is given by the Segre index. By Proposition 3.16, it suffices to show that the local index is equal to the conic index, which is the following theorem.

**Theorem 4.1.** If  $\ell$  is a line on a degree 2n - 1 hypersurface  $X = \mathbb{V}(F)$  in  $\mathbb{P}^{n+1}$ , then  $\operatorname{ind}_{\ell} \sigma_F = \operatorname{con}(X, \ell)$ .

Proof. For any X and  $\ell$ , there exists a rational conic model  $(B, \beta, Q)$  of  $\mathcal{G}(\ell)$  according to Lemma 3.11 and Corollary 3.13. We will prove that  $\operatorname{con}(X, \ell)$  can be expressed as a product of resultants in terms of Q and B, denoted by  $\mathcal{R}(B,Q)$  (see Equation 4.1). Using the coordinate functions of our rational curve  $\mathcal{G}(\ell) = g_{B,\beta}(Q)$ , we will construct a matrix whose determinant only depends on B and Q. We will denote the determinant of this matrix by  $\mathcal{A}(B,Q)$  (Equation 4.6).

So far,  $\mathcal{R}(B,Q)$  and  $\mathcal{A}(B,Q)$  are algebraic maps on  $\operatorname{Conf}_{\binom{n}{2}}(\mathbb{P}^2) \times \mathcal{Q}$ . By applying a projective change of coordinates if necessary, we may assume that  $B \subseteq \mathbb{P}^2$  does not intersect the divisor at infinity, so that  $\mathcal{R}(B,Q)$  and  $\mathcal{A}(B,Q)$  are algebraic maps on  $\operatorname{Conf}_{\binom{n}{2}}(\mathbb{A}^2) \times \mathcal{Q}$ . As  $\mathcal{Q}$  is the space of parameterized conics, we have a parameterization  $[Q_0:Q_1:Q_2]$  of Q. The space of coefficients of a homogeneous degree 2 polynomial in 3 variables is  $\mathbb{A}^3$ , so we may regard  $(Q_0, Q_1, Q_2)$  as a point in  $\mathbb{A}^9$ . As

$$\operatorname{Conf}_{\binom{n}{2}}(\mathbb{A}^2) \subseteq \mathbb{A}^{\binom{n}{2}},$$

we may treat  $\mathcal{R}(B,Q)$  and  $\mathcal{A}(B,Q)$  as algebraic maps on  $\mathbb{A}^{n(n-1)} \times \mathbb{A}^9$ . In order for this change of domain to be well-defined, it suffices to prove that the maps  $\mathcal{R}(B,Q)$  and  $\mathcal{A}(B,Q)$  do not depend (up to squares) on:

- (i) the choice of coordinates on  $\mathbb{P}^2$ ,
- (ii) the choice of parameterization of Q, and
- (iii) the choice of representative of the projective equivalence class  $[Q_0: Q_1: Q_2]$ .

This will be proved in Propositions 4.2 and 4.4. We will then show that

$$\mathcal{R}(B,Q) = \prod_{b \in B} N_{k(b)/k(\ell)} \alpha_b,$$
  
$$\mathcal{A}(B,Q) = \det A_{P_1,\dots,P_n} \text{ (up to squares),}$$

where we are using the notation of Definition 3.15 and Equation 2.1, respectively. If we can prove that  $\mathcal{R}(B,Q) = \mathcal{A}(B,Q)$  up to squares whenever  $\mathcal{A}(B,Q) \neq 0$ , then the claim that

$$\operatorname{con}(X,\ell) = \operatorname{ind}_{\ell} \sigma_F$$

will follow from Definition 3.15 and Equation 2.2.

To prove that  $\mathcal{R}(B,Q) = \mathcal{A}(B,Q)$  up to squares whenever  $\mathcal{A}(B,Q) \neq 0$ , we will show:

- (i) there exists a regular map det  $V_B$  such that if det  $V_B \neq 0$ , then  $\mathcal{A}(B,Q) \neq 0$  (Lemma 4.5),
- (ii)  $(\det \mathbf{V}_B)^{2n} \cdot \mathcal{R}(B,Q)$  and  $\mathcal{A}(B,Q)$  have the same zero locus (Proposition 4.6),
- (iii) there exist infinitely many (B, Q) (over  $\overline{k(\ell)}$ ) such that  $(\det \mathbf{V}_B)^{2n} \cdot \mathcal{R}(B, Q) = \mathcal{A}(B, Q) \neq 0$  (Proposition 4.8), and
- (iv) there exists  $c \in \overline{k(\ell)}$  and  $d \in \mathbb{Z}$  such that  $\mathcal{A}(B,Q) = c \cdot (\det \mathbb{V}_B)^d \cdot \mathcal{R}(B,Q)$ (Proposition 4.11).

We then conclude that  $\mathcal{A}(B,Q) = (\det \mathbb{V}_B)^{2n} \cdot \mathcal{R}(B,Q)$  in Corollary 4.12.

Proof of Theorem 1.2. By Proposition 3.16, the Segre index equals the conic index. By Theorem 4.1, the conic index equals the local degree. This means that the local index equals the Segre index, as claimed.  $\Box$ 

The rest of the paper is devoted to proving the propositions and lemmas referenced in the proof of Theorem 4.1. For the remainder of this section fix a rational conic model  $(B, \beta, Q)$  of  $\mathcal{G}(\ell)$ . Let [Z : X : Y] be coordinates on  $\mathbb{P}^2$  such that  $B \subseteq \{Z \neq 0\}$ . We denote the affine coordinates on  $\{Z \neq 0\}$  by (x, y). Let  $[Q_0 : Q_1 : Q_2]$  be a parameterization of Q (which exists, as Q is a parameterized conic).

4.1. **Defining**  $\mathcal{R}(B,Q)$ . Recall that

$$\operatorname{con}(X,\ell) = \operatorname{Tr}_{k(\ell)/k} \langle \prod_{b \in B} \operatorname{N}_{k(b)/k(\ell)} \alpha_b \rangle.$$

Our goal is to express  $\prod_{b \in B} N_{k(b)/k(\ell)} \alpha_b$  in terms of B and Q, where  $\alpha_b \in k^{\times}$  is such that the fixed points of the involution  $\mu_b$  are defined over  $k(b)(\sqrt{\alpha_b})$ . Let  $b = (b_x, b_y) \in \mathbb{A}^2 = \{Z \neq 0\} \subset \mathbb{P}^2$ . We have

$$\alpha_b = \operatorname{Disc}_{t_0, t_1} \left( \operatorname{Disc}_{u, v} (t_0(Q_1(u, v) - b_x Q_0(u, v)) - t_1(Q_2(u, v) - b_y Q_0(u, v))) \right),$$

which simplifies to

$$\operatorname{Res}(Q_1 - b_x Q_0, Q_2 - b_y Q_0)$$

up to squares. We can now define  $\mathcal{R}(B,Q)$  as

(4.1) 
$$\mathcal{R}(B,Q) = \prod_{b \in B(\overline{k(\ell)})} \operatorname{Res}(Q_1 - b_x Q_0, Q_2 - b_y Q_0).$$

Note that we are taking the product of resultants over the entire geometric locus  $B(k(\ell))$ , which is equal to taking the product of norms of resultants over the locus of closed points B.

**Proposition 4.2.** As an element of  $k(\ell)^{\times}/(k(\ell)^{\times})^2$ , the value  $\mathcal{R}(B,Q)$  does not depend on:

- (i) the choice of coordinates on  $\mathbb{P}^2$ ,
- (ii) the choice of parameterization of Q, and
- (iii) the choice of representative of the projective equivalence class  $[Q_0: Q_1: Q_2]$ .

*Proof.* All three of these statements can be verified computationally, as we now explain.

(i) Any change of coordinates on  $\mathbb{P}^2$  can be represented by (the projective class of) some  $(a_{ij}) = A \in \operatorname{GL}_3(k(\ell))$ . One can compute directly (with your favorite computer algebra system) that after such a coordinate change, we get

(4.2) 
$$(a_{11} + a_{12}b_x + a_{13}b_y)^2 \cdot (\det A)^2 \cdot \operatorname{Res}(Q_2 - b_xQ_1, Q_3 - b_yQ_1),$$

which differs from  $\operatorname{Res}(Q_2 - b_x Q_1, Q_3 - b_y Q_1)$  by a square. Note that even if  $(b_x, b_y)$  is not defined over  $k(\ell)$ , the value  $\mathcal{R}(B, Q)$  is a product over all  $b \in B(\overline{k})$ . Our assumption that B is defined over  $k(\ell)$  implies that the Galois conjugates of Equation 4.2 will also by factors in this product, so that  $\mathcal{R}(B, Q)$  will only change by a square in  $k(\ell)^{\times}$  after our change of coordinates A.

(ii) Choosing a different parametrization of Q is simply precomposing with an automorphism of  $\mathbb{P}^1$ . It suffices to show that a resultant  $\operatorname{Res}(A(z), B(z))$  changes by a square after Möbius transformations when deg  $A = \deg B = 2$ . Applying Möbius transformations to A and B in this case is equivalent to taking new polynomials

$$A'(z) = (cz+d)^2 A\left(\frac{az+b}{cz+d}\right), \qquad B'(z) = (cz+d)^2 A\left(\frac{az+b}{cz+d}\right).$$

Classical properties of resultants imply that

- $\operatorname{Res}(A(z+a), B(z+b)) = \operatorname{Res}(A(z), B(z)),$
- $\operatorname{Res}(A(az), B(az)) = a^{\operatorname{deg} A \cdot \operatorname{deg} B} \operatorname{Res}(A(z), B(z)) = a^4 \operatorname{Res}(A, B)$ , and
- $\operatorname{Res}(z^{\operatorname{deg} A}A(1/z), z^{\operatorname{deg} B}B(1/z)) = (-1)^{\operatorname{deg} A \cdot \operatorname{deg} B} \operatorname{Res}(A(z), B(z)) = \operatorname{Res}(A, B).$

Since Möbius transformations are compositions of translations, invertions and scalar multiplication (which are invariant up to squares), we get the result. In particular,

by computing directly, we can see that

$$\operatorname{Res}(A', B') = \det \begin{pmatrix} a & b \\ c & d \end{pmatrix}^4 \operatorname{Res}(A, B).$$

(iii) For a different choice of projective class, we have  $Q'_i = \lambda Q_i$  for some  $\lambda \in k(\ell)^{\times}$ . Again, it suffices to see that  $\operatorname{Res}(\lambda A, \lambda B) = \operatorname{Res}(A, B)$  up to squares. Indeed, we are simply multiplying the Sylvester matrix by  $\lambda$  and therefore

$$\operatorname{Res}(\lambda A, \lambda B) = \lambda^{\operatorname{deg} A + \operatorname{deg} B} \operatorname{Res}(A, B).$$

In our case this implies that the difference will given by a factor  $\lambda^4$ , which is a square.

Note that the product defining  $\mathcal{R}(B,Q)$  runs over the set of geometric points of B. Each closed point  $b \in B$  consists of a Galois orbit of geometric points, and the product over such an orbit gives the field norm  $N_{k(b)/k(\ell)}$ . In particular, we have

$$\mathcal{R}(B,Q) = \prod_{b \in B} \mathcal{N}_{k(b)/k(\ell)} \alpha_b.$$

4.2. **Defining**  $\mathcal{A}(B,Q)$ . The definition of the regular function  $\mathcal{A}(B,Q)$  is more involved. The strategy is to construct a basis  $\beta'$  in terms of (B,Q). This will allow us to construct a rational curve

$$g_{B,\beta'} \circ [Q_0:Q_1:Q_2]: \mathbb{P}^1 \to \mathbb{P}^n.$$

We will then define the regular function  $\mathcal{A}(B,Q)$  as the determinant of the matrix  $A_{P_1,\ldots,P_n}$  (Equation 2.1), where  $P_1,\ldots,P_n$  are the coordinate functions of  $g_{B,\beta'} \circ [Q_0 : Q_1 : Q_2]$ .

In general,  $\beta'$  need not be the basis coming from our conic model  $(B, \beta, Q)$ . However,  $g_{B,\beta} \circ Q$  and  $g_{B,\beta'} \circ Q$  will be projectively equivalent curves in  $\mathbb{P}^{n-1}$ , and we will prove that the local index of a curve is preserved under projective equivalence (Lemma 4.3).

We will now describe how to construct our basis  $\beta'$ . Let  $B(\overline{k}) = \{b_1, \ldots, b_m\}$ , where  $m = \binom{n}{2}$ . In our chosen affine patch, write  $b_i = (b_{i,x}, b_{i,y})$ . Consider the following interpolation matrix:

(4.3) 
$$\mathbf{V}_B = \begin{pmatrix} 1 & b_{1,x} & b_{1,y} & b_{1,x}^2 & b_{1,x}b_{1,y} & \cdots & b_{1,y}^{n-2} \\ 1 & b_{2,x} & b_{2,y} & b_{2,x}^2 & b_{2,x}b_{2,y} & \cdots & b_{2,y}^{n-2} \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & b_{m,x} & b_{m,y} & b_{m,x}^2 & b_{m,x}b_{m,y} & \cdots & b_{m,y}^{n-2} \end{pmatrix}$$

Note that  $V_B$  is a square  $m \times m$  matrix, since the number of monomials in two variables of degree at most n-2 in two variables is simply  $\sum_{i=1}^{n-1} i = \binom{n}{2}$ . We can think of  $V_B$  as a matrix in the space of polynomials in two variables of degree at most n-2. A solution of the system  $V_B f = w$  is a polynomial f for which  $f(b_{i,x}, b_{i,y}) = w_i$  for all  $b_i \in B$ . If det  $V_B = 0$ , then there is a curve of degree n-2 through all points of B. In this sense,  $V_B$  is a generalization of the Vandermonde matrix (as is also the case for more general interpolation matrices).

In order to get the linear system  $L_B$  of curves of degree n-1 through the points B, we need to add the following n columns to  $V_B$ :

(4.4) 
$$\mathbf{R}_{B} = \begin{pmatrix} b_{1,x}^{n-1} & b_{1,x}^{n-2}b_{1,y} & \cdots & b_{1,y}^{n-1} \\ b_{2,x}^{n-1} & b_{2,x}^{n-2}b_{2,y} & \cdots & b_{2,y}^{n-1} \\ \vdots & \vdots & \ddots & \vdots \\ b_{m,x}^{n-1} & b_{m,x}^{n-2}b_{m,y} & \cdots & b_{m,y}^{n-1} \end{pmatrix}$$

Appending  $\mathbf{R}_B$  to  $\mathbf{V}_B$  yields the  $m \times (m+n)$ -matrix

$$L_B = [V_B \mid R_B].$$

By construction, elements of the kernel of  $L_B$  are exactly the elements of the linear system  $L_B$ . Our next goal is to construct  $\beta'$  as a particular basis of ker  $L_B$ . To begin, let  $V_B^*$  be the adjugate of  $V_B$  and define a block matrix

(4.5) 
$$\mathsf{K}_B := \begin{pmatrix} -\mathsf{V}_B^* \cdot \mathsf{R}_B \\ \det \mathsf{V}_B \cdot \mathsf{I}_n \end{pmatrix},$$

where  $I_n$  denotes the  $n \times n$  identity matrix. Note that  $L_B \cdot K_B = 0$ . If det  $V_B \neq 0$ , the columns of  $K_B$  are linearly independent, in which case the columns of  $K_B$  form a basis for ker  $L_B$ . Let  $\beta'$  be such a basis.

Finally, let Q be the  $(2n+1) \times (m+n)$  matrix representing the linear map

$$k[X, Y, Z]_{(n-1)} \to k[u, v]_{(2n-2)}$$
  
 $f(x, y, z) \mapsto f(Q_0, Q_1, Q_2).$ 

Then  $\mathbb{Q} \cdot \mathbb{K}_B$  is a matrix whose columns give the coefficients of the *n* coordinate polynomials of the parameterized curve  $C = g_{B,\beta'}(Q) \subseteq \mathbb{P}^{n-1}$ . Denote these polynomials by  $P'_1, \ldots, P'_n$ , and let

(4.6) 
$$\mathcal{A}(B,Q) := \det A_{P'_1,\dots,P'_n},$$

where  $A_{P'_1,\ldots,P'_n}$  is the matrix given in Equation 2.1.

Note that if det  $V_B = 0$ , then  $\beta'$  is not necessarily a basis. However, we have det  $V_B \neq 0$  for a general hypersurface in  $\mathbb{P}^n$ . Indeed, if det  $V_B = 0$ , then there exists a degree n-2 curve passing through B. The projective span of this curve is a (2n-4)-secant of dimension n-4 to our hypersurface. As the codimension of this secant is 2, we have infinitely many (2n-4)-secants of codimension 1, which does not occur for general hypersurfaces of degree 2n-1 in  $\mathbb{P}^n$ .

Nevertheless, the matrix  $\mathbf{Q} \cdot \mathbf{K}_{\mathsf{B}}$  can still be computed even when det  $\mathbf{V}_B = 0$  and, therefore, the map  $\mathcal{A}(B,Q)$  can still be defined in this case. In fact, we will prove in Lemma 4.5 that  $\mathcal{A}(B,Q) = 0$  whenever det  $\mathbf{V}_B = 0$ . As previously mentioned, we need to verify that  $\mathcal{A}(B,Q)$  and the determinant of the local index matrix of our Gauß curve  $\mathcal{G}(\ell)$  agree up to squares. This follows from the fact that they differ by a projective change of coordinates:

**Lemma 4.3.** Let  $(B, \beta, Q)$  be a rational conic model of a Gauß curve  $\mathcal{G}(\ell)$ . If  $\beta'$  is another basis of the linear system of degree 2n - 1 curves through B, then the coordinate polynomials  $P_1, \ldots, P_n$  of  $\mathcal{G}(\ell)$  and  $P'_1, \ldots, P'_n$  of  $g_{B,\beta'} \circ Q$  differ by a projective change of coordinates, and

$$\det A_{P_1,\ldots,P_n} = \det A_{P'_1,\ldots,P'_n}$$

up to squares.

Proof. Let  $M = (m_{ij}) \in \operatorname{GL}_n(k(\ell))$  be a matrix such that  $\beta = M \cdot \beta'$ . We then have  $\beta_i = \sum_{j=1}^n m_{ij}\beta'_j$ , and hence composing with our conic Q gives us  $P_i = \sum_{j=1}^n m_{ij}P'_j$ . In particular, the class of M in  $\operatorname{PGL}_n(k(\ell))$  gives us a projective equivalence from  $[P_1 : \ldots : P_n]$  to  $[P'_1 : \ldots : P'_n]$ . Now let

$$\tilde{M} = \begin{pmatrix} m_{11} & 0 & m_{21} & 0 & \cdots & m_{n1} & 0 \\ 0 & m_{11} & 0 & m_{21} & \cdots & 0 & m_{n1} \\ m_{12} & 0 & m_{22} & 0 & \cdots & m_{n2} & 0 \\ 0 & m_{12} & 0 & m_{22} & \cdots & 0 & m_{n2} \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ m_{1n} & 0 & m_{2n} & 0 & \cdots & m_{nn} & 0 \\ 0 & m_{1n} & 0 & m_{2n} & \cdots & 0 & m_{nn} \end{pmatrix}.$$

(In other words,  $\tilde{M} = M \otimes I_2$  is the Kronecker product of M and the  $2 \times 2$  identity matrix.) Note that  $A_{P_1,\ldots,P_n} = A_{P'_1,\ldots,P'_n} \cdot \tilde{M}$ . Moreover, we have det  $\tilde{M} = \det M^2$ , so it follows that det  $A_{P_1,\ldots,P_n} = \det A_{P'_1,\ldots,P'_n}$  up to squares.

Finally, we need to justify that  $\mathcal{A}(B,Q)$  does not depend on the various choices related to picking representatives of B and Q.

**Proposition 4.4.** As an element of  $k(\ell)^{\times}/(k(\ell)^{\times})^2$ , the value  $\mathcal{A}(B,Q)$  does not depend on:

- (i) the choice of coordinates on  $\mathbb{P}^2$ ,
- (ii) the choice of parameterization of Q, and
- (iii) the choice of representative of the projective equivalence class  $[Q_0: Q_1: Q_2]$ .

*Proof.* As with Proposition 4.2, this boils down to some computations.

(i) Let  $\varphi \colon \mathbb{P}^2 \to \mathbb{P}^2$  be a change of coordinates, and let  $B' = \varphi(B)$ . Then  $\varphi$  induces a map  $\tilde{\varphi} \colon L_B \to L_{B'}$  of linear systems given by  $C \mapsto C \circ \varphi^{-1}$ . Pick  $\beta$  be a basis for

 $L_B$ . We then obtain a basis  $\beta'$  for  $L_{B'}$  given by  $\beta'_i = \tilde{\varphi}(\beta_i)$  for each  $\beta_i \in B$  (recall from Lemma 4.3 that the choice of basis does not matter). Now

$$g_{B',\beta'}(\varphi(Q)) = [\beta'_1 \circ \varphi(Q) : \dots : \beta'_n \circ \varphi(Q)]$$
  
=  $[\beta_1 \circ \varphi^{-1} \circ \varphi(Q) : \dots : \beta_n \circ \varphi^{-1} \circ \varphi(Q)]$   
=  $[\beta_1(Q) : \dots : \beta_n(Q)]$   
=  $g_{B,\beta}(Q).$ 

(ii) Let  $[Q_0 : Q_1 : Q_2]$  be a parameterization of Q, and let  $\varphi : \mathbb{P}^1 \to \mathbb{P}^1$  be a Möbius transformation. Then the coordinate polynomials of  $g_{B,\beta}(Q)$  and  $g_{B,\beta}(Q \circ \varphi)$  satisfy

$$\{[P'_1:\ldots:P'_n]\}=\{[P'_1\circ\varphi:\ldots:P'_n\circ\varphi]\},\$$

and hence the coefficient matrices  $A_{P'_1,\ldots,P'_n}$  and  $A_{P'_1\circ\varphi,\ldots,P'_n\circ\varphi}$  differ by  $M \otimes I_2$ , where M is some invertible  $n \times n$  matrix. It follows that the determinants of these two matrices differ by det  $M^2$  (see the proof of Lemma 4.3), which is a square.

(iii) If  $\lambda \in k(\ell)^{\times}$ , then changing  $(Q_0, Q_1, Q_2)$  to  $(\lambda Q_0, \lambda Q_1, \lambda Q_2)$  changes the matrix **Q** by scaling each column by  $\lambda^{n-1}$ . As a result, the polynomials  $P'_1, \ldots, P'_n$  (whose coefficients are given by the columns of  $\mathbf{Q} \cdot \mathbf{K}_B$ ) are sent to polynomials  $\lambda^{n-1}P'_1, \ldots, \lambda^{n-1}P'_n$ . Since  $A_{P'_1,\ldots,P'_n}$  has rank 2n, it follows that

$$\det A_{\lambda^{n-1}P'_1,...,\lambda^{n-1}P'_n} = \lambda^{2n(n-1)} \cdot \det A_{P'_1,...,P'_n},$$

so these two determinants differ by a square.

4.3. Proving  $\mathcal{A}(B,Q) = \mathcal{R}(B,Q)$  up to squares. We have now defined our regular maps  $\mathcal{A}(B,Q)$  and  $\mathcal{R}(B,Q)$  that compute the conic index and local index, respectively. The final step is to show that these two regular maps agree up to squares. Our first goal to this end is to show that  $\mathcal{A}(B,Q)$  and  $(\det V_B)^{2n} \cdot \mathcal{R}(B,Q)$  have the same zero locus.

**Lemma 4.5.** If det  $V_B = 0$ , then  $\mathcal{A}(B, Q) = 0$ .

Proof. If det  $V_B = 0$ , then the bottom n rows of  $K_B$  are 0. In the polynomials corresponding to the columns of  $K_B$ , this means that the coefficients of all monomials that do not contain z are 0. (Recall that here, [x : y : z] are our projective coordinates for  $\mathbb{P}^2$ , and the columns of  $K_B$  give the coefficients of corresponding plane curves.) Hence the n polynomials  $P'_1, \ldots, P'_n$  given by the columns of  $Q \cdot K_B$  have a common factor  $Q_0$ . In particular, this implies that they have a common root  $[u_0 : v_0]$ . Multiplying the matrix  $A_{P'_1,\ldots,P'_n}$  on the left by  $(u_0^{2n-1}, u_0^{2n-2}v_0, \cdots, u_0v_0^{2n-2}, v_0^{2n-1})$ , we get

$$(u_0P'_1(u_0,v_0),v_0P'_1(u_0,v_0),\cdots,u_0P'_n(u_0,v_0),v_0P'_n(u_0,v_0)).$$

Since  $P'_i(u_0, v_0) = 0$  for i = 1, ..., n, we have a nontrivial element in the kernel of the transpose of  $A_{P'_1,...,P'_n}$ , and therefore det  $A_{P'_1,...,P'_n} = \mathcal{A}(B,Q) = 0$ .

The next proposition states that our two regular maps have the same vanishing locus (assuming that det  $V_B \neq 0$ ).

**Proposition 4.6.** The polynomials  $\mathcal{A}(B,Q)$  and  $(\det V_B)^{2n} \cdot \mathcal{R}(B,Q)$  have the same zero locus.

Proof. Note that  $\mathcal{R}(B,Q) = 0$  if and only if  $B \cap Q \neq \emptyset$ . Indeed,  $\mathcal{R}(B,Q) = 0$  if and only if there exists a point  $b \in B(\overline{k})$  for which  $Q_1 - b_x Q_0$  and  $Q_2 - b_y Q_0$  have a common root, which occurs if and only if there is a point  $[u_0 : v_0] \in \mathbb{P}^1(\overline{k})$  for which  $Q_1(u_0, v_0) - b_x Q_0(u_0, v_0) = 0$  and  $Q_2(u_0, v_0) - b_y Q_0(u_0, v_0) = 0$ . Dividing by  $Q_0(u_0, v_0)$ , we find that this condition is equivalent to the existence of  $[u_0 : v_0] \in \mathbb{P}^1_{\overline{k}}$  such that  $Q(u_0, v_0) = [1 : b_x : b_y]$ , which shows that  $p \in Q(\overline{k})$ .

By Lemma 4.5, if det  $V_B = 0$ , then  $\mathcal{A}(B,Q) = 0$ . It thus remains to show that if det  $V_B \neq 0$ , then  $\mathcal{A}(B,Q) = 0$  if and only if  $Q \cap B \neq \emptyset$ . To this end, assume that we have  $b = [1 : b_x : b_y] \in (B \cap Q)(\overline{k})$ . Then the coordinates of the composition map  $g_B \circ Q = \gamma$  have a common root  $[u_0 : v_0]$ , and thus the proof of Lemma 4.5 gives us  $\mathcal{A}(B,Q) = 0$ .

Conversely, if  $\mathcal{A}(B,Q) = 0$ , then there exist linear polynomials  $L_1, \dots, L_n$  (in u, v) such that  $\sum_{i=1}^n L_i P'_i = 0$  by [FK21, Lemma 3.2.2.(3)]. It follows that  $\sum_{i=1}^n L_i \cdot g^{B,\beta'} \circ [Q_0 : Q_1 : Q_2] = 0$ , where  $g^i_{B,\beta'}$  denotes the *i*<sup>th</sup> coordinate of the embedding  $g_{B,\beta'}$ , which corresponds to the basis element  $\beta'_i$ . In particular, the conic Q is contained in the curve  $Y := \sum_{i=1}^n L_i \beta'_i$ , which has degree n and contains the locus B. If  $B \cap Q = \emptyset$ , then all points of B are contained in the components of Y away from Q, so B is contained in a curve of degree n-2. But det  $V_B \neq 0$ , and hence there is no such curve. We thus have  $B \cap Q \neq \emptyset$ , as desired.

Before giving examples of B and Q for which  $\mathcal{A}(B,Q) = (\det V_B)^{2n} \cdot \mathcal{R}(B,Q) \neq 0$ , we need the following lemma about elementary symmetric polynomials.

**Lemma 4.7.** Let  $e_i(X_1, \ldots, X_n)$  denote the *i*<sup>th</sup> elementary symmetric polynomial in *n* variables. Adopt the convention that  $e_0 = 1$ , and  $e_i(X_1, \ldots, X_n) = 0$  if i < 0 or i > n. Then for any  $1 \le j \le n$ , we have

$$e_{i+1}(X_1,\ldots,X_n) = \sum_{z=0}^{i+1} e_z(X_1,\ldots,X_j)e_{i-z+1}(X_{j+1},\ldots,X_n).$$

*Proof.* Let

$$E_z := e_z(X_1,\ldots,X_j)e_{i-z+1}(X_{j+1},\ldots,X_n).$$

Each of the summands  $E_z$  consists of a sum of distinct monomials of degree i + 1, each with coefficient 1. Moreover, given a monomial summand  $M := X_1^{c_1} \cdots X_n^{c_n}$  of  $e_{i+1}(X_1, \ldots, X_n)$  (so  $c_{\ell} \in \{0, 1\}$  and  $\sum_{\ell=1}^n c_{\ell} = i + 1$ ), there is at most one z such that M is a summand of  $E_z$ . Indeed, any monomial summand  $X_1^{d_1} \cdots X_n^{d_n}$  of  $E_z$  must satisfy  $\sum_{\ell=1}^j d_\ell = z$ .

It therefore suffices to show that the numbers of monomial summands in  $\sum_{z=0}^{i+1} E_z$  and  $e_{i+1}(X_1, \ldots, X_n)$  are equal. The latter number is  $\binom{n}{i+1}$ , essentially by definition of the elementary symmetric polynomials. Similarly, the number of summands in  $e_z(X_1, \ldots, X_j)$ 

and  $e_{i-z+1}(X_{j+1},\ldots,X_n)$  are given by  $\binom{j}{z}$  and  $\binom{n-j}{i-z+1}$ , respectively. Since the indeterminates for these two elementary symmetric polynomials are disjoint, the product  $E_z$  consists of  $\binom{j}{z}\binom{n-j}{i-z+1}$  distinct monomials. Summing up to count the distinct monomials of  $\sum_{z=0}^{i+1} E_z$ , we find that it suffices to prove

$$\binom{n}{i+1} = \sum_{z=0}^{i+1} \binom{j}{z} \binom{n-j}{i-z+1}.$$

This follows from the binomial theorem by comparing the coefficient of  $x^{i+1}$  on both sides of the equation

$$(x+1)^n = (x+1)^j (x+1)^{n-j}.$$

Equivalently (but perhaps more illustratively), repeated application of the standard identity  $\binom{n-m}{i-z} + \binom{n-m}{i-z+1} = \binom{n-m+1}{i-z+1}$  implies that the following rows all have the same sum:

$$\binom{n}{i+1} + \binom{n-1}{i}$$

$$\binom{n-1}{i+1} + \binom{n-1}{i}$$

$$\binom{n-2}{i+1} + 2\binom{n-2}{i} + \binom{n-2}{i-1}$$

$$\binom{n-3}{i+1} + 3\binom{n-3}{i} + 3\binom{n-3}{i-1} + \binom{n-3}{i-2}$$

$$\binom{n-4}{i+1} + 4\binom{n-4}{i} + 6\binom{n-4}{i-1} + 4\binom{n-4}{i-2} + \binom{n-4}{i-3}. \square$$

**Proposition 4.8.** Let  $a_1, \ldots, a_n \in \overline{k(\ell)}$  be distinct elements. Define (4.7)  $B := \{ [1:a_i:a_j] \in \mathbb{P}^2 \mid 1 \le i < j \le n \},\$ 

and  $Q_1 = Q_2 = u^2$  and  $Q_0 = v^2$ . For these choices of B and Q, we have  $\mathcal{A}(B,Q) = \det(\mathbb{V}_B)^{2n} \cdot \mathcal{R}(B,Q) \neq 0$ .

*Proof.* We will prove this in four steps.

Step 1:  $\mathcal{R}(B,Q) = \prod_{i < j} (a_i - a_j)^2$ . We prove this step directly. For each point  $[1 : a_i : a_j] \in B$ , we have  $Q_1 - a_i Q_0 = u^2 - a_i v^2$  and  $Q_2 - a_j Q_0 = u^2 - a_j v^2$ . Thus

$$\mathcal{R}(B,Q) = \prod_{1 \le i < j \le n} \operatorname{Res}(u^2 - a_i v^2, u^2 - a_j v^2)$$
$$= \prod_{1 \le i < j \le n} (a_i - a_j)^2.$$

Step 2: det  $V_B \neq 0$ . We prove this step by contradiction. If det  $V_B = 0$ , then there exists a non-zero polynomial f(x, y) of degree n - 2 such that  $f(a_i, a_j) = 0$  for all  $1 \leq i < j \leq n$ . If we fix  $x = a_1$ , we get a polynomial  $f(a_1, y)$  of degree at most n - 2 with n - 1 distinct roots  $(y = a_2, \ldots, a_n)$ . It follows that  $f(a_1, y) = 0$ , so  $x - a_1$  is a factor of f(x, y). Let  $f_1(x, y) = \frac{f(x, y)}{x - a_1}$ . Then  $f_1(a_1, y)$  is a polynomial of degree at most n-3 with n-2 distinct roots, and hence  $x-a_2$  is a factor of  $f_1(x, y)$ . Repeating this process, we find that  $x-a_i$  is a factor of f(x, y) for all i, and hence f(x, y) = 0 (as the deg(f) < n-1). This contradicts the fact that f was a non-zero polynomial, so we find that det  $V_B \neq 0$ .

Step 3: With our particular choice of  $[Q_0 : Q_1 : Q_2]$  and B, we have

$$A_{P_1',\dots,P_n'} = \det \mathbb{V}_B \cdot \begin{pmatrix} 1 & 0 & \cdots & 1 & 0 \\ 0 & 1 & \cdots & 0 & 1 \\ p_{2n-4,1} & 0 & \cdots & p_{2n-4,n} & 0 \\ 0 & p_{2n-4,1} & \cdots & 0 & p_{2n-4,n} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ p_{0,1} & 0 & \cdots & p_{0,n} & 0 \\ 0 & p_{0,1} & \cdots & 0 & p_{0,n} \end{pmatrix}$$

where

$$p_{j,i} = \begin{cases} (-1)^{\ell-1} e_{\ell}(a_1, \dots, \hat{a}_{n-i+1}, \dots, a_n) & j = 2n - 2 - 2\ell \\ 0 & j \text{ odd} \end{cases}$$

and  $e_{\ell}(X_1, \ldots, X_{n-1})$  is the  $\ell^{\text{th}}$  elementary symmetric polynomial in n-1 variables. (Here, we use the conventions that  $e_0 = 1$  and  $\hat{a}_i$  means that  $a_i$  is omitted.)

Proving this formula is the lengthiest of our four steps. To begin, we need to construct our basis  $\beta'$  from the columns of

$$\mathbf{K}_B = \begin{pmatrix} -\mathbf{V}_B^* \cdot \mathbf{R}_B \\ \det \mathbf{V}_B \cdot \mathbf{I}_n \end{pmatrix}$$

We can then compute  $(p_{j,i}) = \mathbb{Q} \cdot \mathbb{K}_B$ , where  $\mathbb{Q} : k[X, Y, Z]_{(n-1)} \to k[u, v]_{(2n-2)}$  is the linear transformation given by

$$\mathsf{Q}(X^i Y^j Z^{n-1-i-j}) = u^{2(i+j)} v^{2(n-1-i-j)}.$$

Since det  $V_B \neq 0$ , we have  $V_B^* = \det V_B \cdot V_B^{-1}$ , and hence

$$\mathbf{K}_B = \det \mathbf{V}_B \cdot \begin{pmatrix} -\mathbf{V}_B^{-1} \cdot \mathbf{R}_B \\ \mathbf{I}_n \end{pmatrix}.$$

We may therefore compute the columns of  $K_B$  by computing the columns of

$$\mathbf{K}_B' := \begin{pmatrix} -\mathbf{V}_B^{-1} \cdot \mathbf{R}_B \\ \mathbf{I}_n \end{pmatrix}$$

From now on, we interpret the columns of  $K'_B$  as polynomials in x, y (rather than homogeneous polynomials in x, y, z) by setting z = 1. As identity matrices are easy enough to understand, we turn our attention to  $V_B^{-1} \cdot R_B$ . For each column  $w_j$  of  $R_B$  (see Equation 4.4), we need to solve the system  $V_B \cdot f_j = w_j$  for the column  $f_j$ . That is, we need to find a degree n - 2 polynomial  $f_j \in k(\ell)[x, y]$ (which we conflate with its column of coefficients) such that  $f_j(b_{i,x}, b_{i,y}) = w_{i,j}$  for all  $[1: b_{i,x}: b_{i,y}] \in B$ , where  $w_{i,j}$  denotes the  $i^{\text{th}}$  row of  $w_j$ . It follows that in order to solve  $f_j(b_{i,x}, b_{i,y}) = w_{i,j}$ , it suffices to find a polynomial  $f_j$  such that

$$g_j(x,y) := f_j(x,y) - x^{n-j}y^{j-1}$$

satisfies  $g_j(b_{i,x}, b_{i,y}) = 0$  for all  $1 \le i \le {n \choose 2}$ . Let  $r_{s,t}^j \in k(\ell)$  be such that

$$f_j(x,y) = \sum_{s=0}^{n-2} \left( \sum_{t=0}^s r_{s,t}^j y^t \right) x^{n-2-s}.$$

Let  $r_s^j(y) = \sum_{t=0}^s r_{s,t}^j y^t$ , so that  $f_j(x,y) = \sum_{s=0}^{n-2} r_s^j(y) x^{n-2-s}$ .

We need to compute  $r_{s,t}^j$  such that  $g_j(b_{i,x}, b_{i,y}) = 0$  for all *i*. We will compute  $r_{s,t}^1$  and  $r_{s,t}^2$  to illustrate our general approach, after which we will compute  $r_{s,t}^j$  for all *j*. Recall that for our choice of *B*, we have

$$(b_{1,x}, b_{1,y}) = (a_1, a_2),$$
  

$$(b_{n-1,x}, b_{n-1,y}) = (a_1, a_n),$$
  

$$(b_{n,x}, b_{n,y}) = (a_2, a_3),$$
  

$$(b_{2n-3,x}, b_{2n-3,y}) = (a_2, a_n),$$
  

$$\vdots$$
  

$$(b_{\binom{n}{2},x}, b_{\binom{n}{2},y}) = (a_{n-1}, a_n).$$

For j = 1, we have  $g_1(x, y) = -x^{n-1} + f_1(x, y)$ . If  $g_1(b_{i,x}, b_{i,y}) = 0$  for all i, then  $g_1(x, a_n)$  is a degree n - 1 polynomial in x with roots  $a_1, \ldots, a_{n-1}$ . By Vieta's formulas, it follows that  $r_{0,0}^1 = a_1 + \ldots + a_{n-1}$ . Note that for any fixed  $y_0$ , the roots of  $g_1(x, y_0)$  must also sum to  $a_1 + \ldots + a_{n-1}$ . Thus the roots of  $g_1(x, a_{n-1})$  are given by  $a_1, \ldots, a_{n-2}$  (by our definition of B) and also  $a_{n-1}$  (since the roots must sum to  $a_1 + \ldots + a_{n-1}$ ). Vieta's formulas now give us

$$-\sum_{1 \le i < j \le n-1} a_i a_j = r_{1,0}^1 + r_{1,1}^1 a_n$$
$$= r_{1,0}^1 + r_{1,1}^1 a_{n-1}$$

so  $r_{1,1}^1 = 0$  and  $r_{1,0}^1 = -\sum_{i < j} a_i a_j$ . Continuing this process for  $y = a_{n-2}, a_{n-3}, \ldots$ , we conclude that

$$r_s^1(y) = (-1)^s e_{s+1}(a_1, \dots, a_{n-1}),$$

where  $e_{s+1}$  denotes the  $(s+1)^{st}$  elementary symmetric polynomial in n-1 variables.

For j = 2, we have  $g_2(x, y) = (r_{0,0}^2 - y)x^{n-2} + \sum_{s=1}^{n-2} r_s^2(y)x^{n-2-s}$ . As with j = 1, we will consider  $g_2(x, a_n)$ . We then have n - 1 roots  $(a_1, \ldots, a_{n-1})$  of this degree n-2 polynomial, so  $g_2(x, a_n)$  must be identically zero. It follows that  $r_s^2(a_n) = 0$ 

for all s, so each of these polynomials is divisible by  $y - a_n$ . Let  $\bar{r}_s^2(y) = \frac{r_s^2(y)}{y - a_n}$ . Factoring, we have

$$g_2(x,y) = (y-a_n) \left( -x^{n-2} + \sum_{s=1}^{n-2} \bar{r}_s^2(y) x^{n-2-s} \right)$$

Now since  $a_i \neq a_n$  for i < n, the assumption that  $g_2(a_i, a_n) = 0$  implies that the polynomial

$$-x^{n-2} + \sum_{s=1}^{n-2} \bar{r}_s^2(a_{n-1})x^{n-2-s}$$

has roots  $a_1, \ldots, a_{n-2}$ . Repeating our arguments from the j = 1 case, we find that  $\bar{r}_s^2(y) = (-1)^s e_s(a_1, \ldots, a_{n-2})$ , where we now have the  $s^{\text{th}}$  elementary symmetric polynomial in n-2 variables (rather than the  $(s+1)^{\text{st}}$  elementary symmetric polynomial in n-1 variables). In particular, we have  $r_s^2 = (-1)^s (y-a_n) e_s(a_1, \ldots, a_{n-2})$  and hence

$$r_{s,t}^{2} = \begin{cases} (-1)^{s+1} a_{n} e_{s}(a_{1}, \dots, a_{n-2}), & t = 0, \\ (-1)^{s} e_{s}(a_{1}, \dots, a_{n-2}) & t = 1, \\ 0 & t \ge 2. \end{cases}$$

For j > 2, we follow the same steps as in the j = 2 case. We find that  $a_1, \ldots, a_{n-1}$  are roots of  $g_j(x, a_n)$ , so  $g_j(x, a_n) = 0$ . Thus  $r_{0,0}^j = 0$ , so  $g_j(x, a_n)$  has degree n-3 in x. This implies that  $g_j(x, a_{n-1}) = 0$ , which means that the linear polynomial  $r_1^j(y)$  has distinct roots  $a_n$  and  $a_{n-1}$  and must therefore be identically zero, and likewise for the polynomials  $r_s^j(y)$  with s < j - 2 upon repeating this process. We conclude that

$$g_j(x,y) = \prod_{i=0}^{j-2} (y - a_{n-i}) \cdot \left( -x^{n-j} + \sum_{s=j-1}^{n-2} \bar{r}_s^j(y) x^{n-2-s} \right),$$

where

$$\bar{r}_s^j(y) \cdot \prod_{i=0}^{j-2} (y - a_{n-i}) = r_s^j(y).$$

(4.8)

Since  $g_j(x, a_{n-j+1}) = 0$  and  $\prod_{i=0}^{j-2} (a_{n-j+1} - a_{n-i}) \neq 0$ , it follows that the degree n-j polynomial

$$-x^{n-j} + \sum_{s=j-1}^{n-2} \bar{r}_s^j(a_{n-j+1})x^{n-2-s}$$

has roots  $a_1, \ldots, a_{n-j}$ . As in the j = 2 case, we find that

$$\bar{r}_s^j(y) = (-1)^s e_{s-j+2}(a_1, \dots, a_{n-j})$$

It now follows from Equation 4.8 and Vieta's formulas that

(4.9) 
$$r_{s,t}^{j} = (-1)^{s+t+1} e_{s-j+2}(a_1, \dots, a_{n-j}) e_{j-1-t}(a_{n-j+2}, \dots, a_n),$$

where we use the convention that  $e_0 = 1$  and  $e_m = 0$  for m < 0.

We have solved for the columns of  $V_B^{-1} \cdot R_B$ . When considering the columns of  $K'_B$ , we must multiply the columns of  $V_B^{-1} \cdot R_B$  by -1. After composing with  $[Q_0 : Q_1 : Q_2]$  (i.e. multiplying by  $\mathbb{Q}$ ), the monomials corresponding to the coefficients of the lower block  $\mathbf{I}_n$  are all  $u^{2n-2}$  for our choice of conic. It follows that

$$\mathbf{Q} \cdot \mathbf{K}'_B = \begin{pmatrix} 1 & 1 & \cdots & 1 \\ 0 & 0 & \cdots & 0 \\ -\sum_{s=0}^{n-2} r_{s,s}^1 & -\sum_{s=0}^{n-2} r_{s,s}^2 & \cdots & -\sum_{s=0}^{n-2} r_{s,s}^n \\ 0 & 0 & \cdots & 0 \\ -\sum_{s=0}^{n-3} r_{s+1,s}^1 & -\sum_{s=0}^{n-3} r_{s+1,s}^2 & \cdots & -\sum_{s=0}^{n-3} r_{s+1,s}^n \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \\ -(r_{n-3,0}^1 + r_{n-2,1}^1) & -(r_{n-3,0}^2 + r_{n-2,1}^2) & \cdots & -(r_{n-3,0}^n + r_{n-2,1}^n) \\ 0 & 0 & \cdots & 0 \\ -r_{n-2,0}^1 & -r_{n-2,0}^2 & \cdots & -r_{n-2,0}^n \end{pmatrix}.$$

To complete Step 3, it remains to show that

$$-\sum_{s=0}^{n-2-i} r_{s+i,s}^{j} = (-1)^{i} e_{i+1}(a_1, \dots, \hat{a}_j, \dots, a_n).$$

By Equation 4.9, we have

$$r_{s+i,s}^{j} = (-1)^{i+1} e_{s+i-j+2}(a_1, \dots, a_{n-j}) e_{j-1-s}(a_{n-j+2}, \dots, a_n).$$

Note that  $r_{s+i,s}^j = 0$  when s < j - i - 2 or s > j - 1, so

$$-\sum_{s=0}^{n-2-i} r_{s+i,s}^j = -\sum_{s=j-i-2}^{j-1} r_{s+i,s}^j.$$

We will re-index by setting z = s - (j - i - 2), so that

$$-\sum_{s=j-i-2}^{j-1} r_{s+i,s}^{j} = -\sum_{z=0}^{i+1} r_{z+j-2,z+j-i-2}^{j}$$
$$= (-1)^{i+2} \sum_{z=0}^{i+1} e_{z}(a_{1},\ldots,a_{n-j})e_{i-z+1}(a_{n-j+2},\ldots,a_{n}).$$

It follows from Lemma 4.7 that

$$\sum_{z=0}^{i+1} e_z(a_1, \dots, a_{n-j}) e_{i-z+1}(a_{n-j+2}, \dots, a_n) = e_{i+1}(a_1, \dots, \hat{a}_{n-j+1}, \dots, a_n),$$

as required.

Step 4: 
$$\mathcal{A}(B,Q) = (\det \mathbb{V}_B)^{2n} \cdot \prod_{i < j} (a_i - a_j)^2$$
. By Step 3, we have  
 $\mathcal{A}(B,Q) = \det A_{P'_1,\dots,P'_n}$ 

$$= (\det \mathbb{V}_B)^{2n} \cdot \det \begin{pmatrix} 1 & 0 & \cdots & 1 & 0 \\ 0 & 1 & \cdots & 0 & 1 \\ p_{2n-4,1} & 0 & \cdots & p_{2n-4,n} & 0 \\ 0 & p_{2n-4,1} & \cdots & 0 & p_{2n-4,n} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ p_{0,1} & 0 & \cdots & p_{0,n} & 0 \\ 0 & p_{0,1} & \cdots & 0 & p_{0,n} \end{pmatrix},$$

so it suffices to prove that the latter matrix (which we call  $\tilde{N}$ ) has determinant  $\prod_{i < j} (a_i - a_j)^2$ . Note that  $\tilde{N} = N \otimes I_2$ , where

$$N := \begin{pmatrix} 1 & 1 & \cdots & 1 \\ p_{2n-4,1} & p_{2n-4,2} & \cdots & p_{2n-4,n} \\ \vdots & \vdots & \ddots & \vdots \\ p_{0,1} & p_{0,2} & \cdots & p_{0,n} \end{pmatrix},$$

so det  $\tilde{N} = \det N^2$ . It therefore suffices to prove that

$$\det N = \pm \prod_{1 \le i < j \le n} (a_i - a_j).$$

which we will do by induction on n. Note that  $p_{i,j}$  is a polynomial in n-1 variables, even though this dependence on n is not reflected in the notation. To make this dependence more explicit, let  $e_{\ell}(\boldsymbol{a}_{i,n}) = e_{\ell}(a_1, \ldots, \hat{a}_i, \ldots, a_n)$  and

$$N_{n} = \begin{pmatrix} 1 & 1 & \cdots & 1 \\ e_{1}(\boldsymbol{a}_{n,n}) & e_{1}(\boldsymbol{a}_{n-1,n}) & \cdots & e_{1}(\boldsymbol{a}_{1,n}) \\ \vdots & \vdots & \ddots & \vdots \\ (-1)^{n-2}e_{n-1}(\boldsymbol{a}_{n,n}) & (-1)^{n-2}e_{n-1}(\boldsymbol{a}_{n-1,n}) & \cdots & (-1)^{n-2}e_{n-1}(\boldsymbol{a}_{1,n}) \end{pmatrix}$$

If n = 2, then we have

$$N_2 = \begin{pmatrix} 1 & 1 \\ a_2 & a_1 \end{pmatrix},$$

and thus det  $N_2 = a_1 - a_2$ , as desired. Next, assume that

$$\det N_n = \pm \prod_{1 \le i < j \le n} (a_i - a_j)^2$$

In order to compute det  $N_{n+1}$ , subtract the first column of  $N_{n+1}$  from the last n columns of  $N_{n+1}$ . Since

$$e_{\ell}(\boldsymbol{a}_{i,n+1}) - e_{\ell}(\boldsymbol{a}_{n+1,n+1}) = (a_{n+1} - a_i)e_{\ell-1}(\boldsymbol{a}_{i,n}),$$

it follows that det  $N_{n+1}$  is equal to

$$\det \begin{pmatrix} 1 & 0 & \cdots & 0 \\ e_1(\boldsymbol{a}_{n+1,n+1}) & (a_{n+1}-a_n)e_0(\boldsymbol{a}_{n,n}) & \cdots & (a_{n+1}-a_1)e_0(\boldsymbol{a}_{1,n}) \\ \vdots & \vdots & \ddots & \vdots \\ (-1)^{n-1}e_n(\boldsymbol{a}_{n+1,n+1}) & (-1)^{n-1}(a_{n+1}-a_n)e_{n-1}(\boldsymbol{a}_{n,n}) & \cdots & (-1)^{n-1}(a_{n+1}-a_1)e_{n-1}(\boldsymbol{a}_{1,n}) \end{pmatrix}.$$

After factoring  $a_{n+1} - a_i$  out of the  $(n+2-i)^{\text{th}}$  column for i = 1, ..., n and a -1 out of last n-1 rows, the resulting matrix has the form

$$\begin{pmatrix}
1 & 0 & \cdots & 0 \\
* & & \\
\vdots & N_n \\
* & & 
\end{pmatrix}.$$

It follows that

$$\det N_{n+1} = (-1)^{n-1} \prod_{i=1}^{n} (a_{n+1} - a_i) \cdot \det N_n$$
$$= -\prod_{i=1}^{n} (a_i - a_{n+1}) \cdot \det N_n.$$

We thus have

$$\det N_{n+1} = \mp \prod_{1 \le i < j \le n+1} (a_i - a_j)$$

which completes the proof by induction.

Our final goal is to show that  $\mathcal{A}(B,Q)$  and  $(\det V_B)^{2n} \cdot \mathcal{R}(B,Q)$  differ by a constant, which we will then show to be one. Before proving this, we need the following lemmas.

**Lemma 4.9.** As a polynomial on  $\mathbb{A}^{2\binom{n}{2}} \times \mathbb{A}^9$  (in the coordinates of B and coefficients of Q), the polynomials

$$\operatorname{Res}(Q_1 - b_x Q_0, Q_2 - b_y Q_0)$$

are irreducible over  $\overline{k(\ell)}$ .

*Proof.* We can directly compute these resultants using the following Sage code.

R. <b\_x, b\_y, q\_00, q\_01, q\_02, \\
 q\_10, q\_11, q\_12, q\_20, q\_21, q\_22> = QQ[];
S. <u> = R[];
Q\_0 = q\_00 + q\_01\*u + q\_02\*u^2;
Q\_1 = q\_10 + q\_11\*u + q\_12\*u^2;
Q\_2 = q\_20 + q\_21\*u + q\_22\*u^2;
f = Q\_1 - b\_x\*Q\_0;
g = Q\_2 - b\_y\*Q\_0;
print(f.resultant(g))

Let  $T = k(\ell)[q_{00}, \ldots, q_{22}, b_x]$  with fraction field K. It is then straightforward to check that  $h(b_y) := \operatorname{Res}(Q_1 - b_x Q_0, Q_2 - b_y Q_0) \in T[b_y]$  is a quadratic polynomial. Moreover,

if we write  $h(b_y) = h_2 b_y^2 + h_1 b_y + h_0$ , we can compute that  $gcd(h_0, h_1, h_2) = 1$  (either with Sage, or by noting that no terms of  $h_2$  are divisible by  $b_x$ , but some terms of  $h_0$ and  $h_1$  are divisible by  $b_x$ ). Using the quadratic formula (and assuming char  $k \neq 2$ ), it is straightforward to check that  $h(b_y)$  is irreducible over  $K[b_y]$ . It now follows from Gauß's lemma that  $h(b_y)$  is irreducible over  $T[b_y]$ , as desired.

**Lemma 4.10.** The polynomial det  $V_B$  in the coordinates  $b_{i,x}$  and  $b_{i,y}$  is irreducible over  $\overline{k(\ell)}$ .

*Proof.* Recall that we have assumed  $n \ge 3$  (Remark 3.1), so  $\binom{n}{2} \ge 3$ . In the notation of [DT09, Theorem 1.5], we have  $N = \binom{n}{2}$ ,

$$(\gamma_{1_1}, \gamma_{2_1}, \dots, \gamma_{N_1}) = (0, 1, 0, 2, 1, 0, \dots, n-2, n-3, \dots, 0), (\gamma_{1_2}, \gamma_{2_2}, \dots, \gamma_{N_2}) = (0, 0, 1, 0, 1, 2, \dots, 0, 1, \dots, n-2),$$

and  $\overline{\gamma} = (0,0)$ . Since  $\gamma_2 = (1,0)$ , the largest natural number  $d_{\Gamma}$  such that  $\frac{1}{d_{\Gamma}}(\gamma_2 - \overline{\gamma}) = (\frac{1}{d_{\Gamma}}, 0)$  is an element of  $\mathbb{N}^2$  is 1. Finally, since  $\gamma_2 = (1,0)$  and  $\gamma_3 = (0,1)$ , it follows that dim  $\mathcal{L}_{\Gamma} = 2$ . Therefore the assumptions of [DT09, Theorem 1.5] hold, which implies that the interpolation determinant det  $\mathbb{V}_B$  is irreducible over any algebraically closed field.  $\Box$ 

We can now show that  $\mathcal{A}(B,Q)$  and  $\mathcal{R}(B,Q)$  differ by  $c \cdot (\det \mathbf{V}_B)^d$  for some constant c and some integer d.

**Proposition 4.11.** As polynomials on  $\mathbb{A}^{\binom{n}{2}} \times \mathbb{A}^9$ , we have  $\mathcal{A}(B,Q) = c \cdot \det(\mathbb{V}_B)^d \cdot \mathcal{R}(B,Q)$  for some  $c \in \overline{k(\ell)}$  and  $d \in \mathbb{Z}$ .

*Proof.* Since  $\mathcal{A}$  and det $(V_B)^{2n} \cdot \mathcal{R}$  have the same zero locus (Proposition 4.6), Hilbert's Nullstellensatz implies that radical ideals generated by the two polynomials are the same. Therefore

(4.10) 
$$\sqrt{(\mathcal{A})} = \sqrt{(\det \mathbf{V}_B)} \cap \sqrt{(\mathcal{R})}.$$

Since  $\mathcal{R}$  is the product of distinct irreducible polynomials (over  $\overline{k(\ell)}$ ) and det  $V_B$  is irreducible (over  $\overline{k(\ell)}$ ) by Lemmas 4.9 and 4.10, we conclude that the ideals ( $\mathcal{R}$ ) and (det  $V_B$ ) are radical. This implies that

(4.11) 
$$\mathcal{A} = c \cdot \det(\mathbb{V}_B)^d \cdot \prod_{b \in B(\overline{k(\ell)})} \operatorname{Res}(Q_1 - b_x Q_0, Q_2 - b_y Q_0)^{N_i}$$

for some integers d > 0 and  $N_i > 0$  and some  $c \in \overline{k(\ell)}$ .

We finish by showing that  $N_i = 1$  for  $i = 1, ..., \binom{n}{2}$ . Consider the value of the polynomial  $\mathcal{A}$  for Q and  $\lambda Q$  for some  $\lambda \in \overline{k(\ell)}$ . By Proposition 4.2 (iii), we have  $\operatorname{Res}(\lambda Q_1 - \lambda b_x Q_0, \lambda Q_2 - \lambda b_y Q_0) = \lambda^4 \operatorname{Res}(Q_1 - b_x Q_0, Q_2 - b_y Q_0)$ . Similarly, by Proposition 4.4 (iii),

we have  $\mathcal{A}(\lambda Q, B) = \lambda^{2n(n-1)} \mathcal{A}(Q, B)$ . Comparing the multiplicity of  $\lambda$  in Equation 4.11, we get

$$4\sum_{i=1}^{\binom{n}{2}} N_i = 2n(n-1),$$

which implies that  $\sum_{i} N_i = {n \choose 2}$ . Since  $N_i > 0$  for each *i*, we have  $N_i = 1$  for all *i*.  $\Box$ 

Finally, we conclude by using Proposition 4.8 to show that c = 1 and d = 2n in Equation 4.11.

**Corollary 4.12.** We have that c = 1 and d = 2n in equation 4.11. Therefore, up to squares,  $\mathcal{A}(B,Q) = \mathcal{R}(B,Q)$ , which implies that the conic index is equal to the local index, as desired.

*Proof.* Proposition 4.8 and Equation 4.11 imply that

 $c \cdot (\det \mathbf{V}_B)^d = (\det \mathbf{V}_B)^{2n}$ 

for any B of the form given in Equation (4.7). It follows that  $(\det V_B)^{2n-d}$  is constant for any such B, which is true only if 2n - d = 0. Consequently, c = 1.

## References

- [BBM<sup>+</sup>21] Thomas Brazelton, Robert Burklund, Stephen McKean, Michael Montoro, and Morgan Opie. The trace of the local A<sup>1</sup>-degree. *Homology Homotopy Appl.*, 23(1):243–255, 2021.
- [BW23] Tom Bachmann and Kirsten Wickelgren. Euler classes: six-functors formalism, dualities, integrality and linear subspaces of complete intersections. J. Inst. Math. Jussieu, 22(2):681– 746, 2023.
- [Cay49] Arthur Cayley. On the triple tangent planes to a surface of the third order. Cambridge and Dublin Mathematical Journal, 4:252–260, 1849.
- [CSD07] F. Chen, L. Shen, and J. Deng. Implicitization and parametrization of quadratic and cubic surfaces by μ-bases. *Computing*, 79(2-4):131–142, 2007.
- [DT09] Carlos D'Andrea and Luis Felipe Tabera. Tropicalization and irreducibility of generalized Vandermonde determinants. *Proc. Amer. Math. Soc.*, 137(11):3647–3656, 2009.
- [Dye97] R. H. Dye. Space sextic curves with six bitangents, and some geometry of the diagonal cubic surface. Proc. Edinb. Math. Soc., II. Ser., 40(1):85–97, 1997.
- [EH16] David Eisenbud and Joe Harris. 3264 and all that. A second course in algebraic geometry. Cambridge: Cambridge University Press, 2016.
- [FK12] Sergey Finashin and Viatcheslav Kharlamov. Abundance of Real Lines on Real Projective Hypersurfaces. International Mathematics Research Notices, 2013(16):3639–3646, 06 2012.
- [FK21] Sergey Finashin and Viatcheslav Kharlamov. Segre indices and Welschinger weights as options for invariant count of real lines. Int. Math. Res. Not., 2021(6):4051–4078, 2021.
- [Kle74] Steven L. Kleiman. The transversality of a general translate. *Compositio Math.*, 28:287–297, 1974.
- [KW19] Jesse Leo Kass and Kirsten Wickelgren. The class of Eisenbud-Khimshiashvili-Levine is the local A<sup>1</sup>-Brouwer degree. Duke Math. J., 168(3):429–469, 2019.
- [KW21] Jesse Leo Kass and Kirsten Wickelgren. An arithmetic count of the lines on a smooth cubic surface. Compos. Math., 157(4):677–709, 2021.
- [Lev19] Marc Levine. Motivic Euler characteristics and Witt-valued characteristic classes. Nagoya Math. J., 236:251–310, 2019.
- [McK22] Stephen McKean. Circles of Apollonius two ways. arXiv:2210.13288, 2022.

- [McK25] Stephen McKean. Rational lines on smooth cubic surfaces. arXiv:2101.08217, 2025.
- [OT14] Christian Okonek and Andrei Teleman. Intrinsic signs and lower bounds in real algebraic geometry. Journal für die reine und angewandte Mathematik (Crelles Journal), 2014(688):219– 241, 2014.
- [Pau22] Sabrina Pauli. Quadratic types and the dynamic Euler number of lines on a quintic threefold. Adv. Math., 405:Paper No. 108508, 37, 2022.
- [Sch58] Ludwig Schläfli. An attempt to determine the twenty-seven lines upon a surface of the third order, and to divide such surfaces into species in reference to the reality of the lines upon the surface. Quart. J. Pure Appl. Math., (2):110–120, 1858.
- [Seg42] B. Segre. The Non-singular Cubic Surfaces: A New Method of Investigation with Special Reference to Questions of Reality. Clarendon Press, 1942.
- [Sol06] Jake P. Solomon. Intersection theory on the moduli space of holomorphic curves with Lagrangian boundary conditions. Preprint, arXiv:math/0606429 [math.SG] (2006), 2006.

Institute de Mathématiques de Jussieu, Sorbonne Université

Email address: espreafico-guelerman@imj-prg.fr

DEPARTMENT OF MATHEMATICS, BRIGHAM YOUNG UNIVERSITY

Email address: mckean@math.byu.edu

 $\mathit{URL}: \mathtt{shmckean.github.io}$ 

FACHBEREICH MATHEMATIK, TU DARMSTADT

Email address: pauli@mathematik.tu-darmstadt.de

URL: homepage.sabrinapauli.com